## Medicare STARS Program

The Centers for Medicare and Medicaid Services (CMS) works with Medicare Advantage Plans like GlobalHealth (HMO) and Generations Healthcare HMO to improve the quality, performance and cost effectiveness of services provided to Medicare beneficiaries. The Medicare STARS program determines how well plans and providers perform across a section of quality measures using information from member satisfaction (both plan and providers), health outcomes and plan operations. A plan can get a rating from one to five stars. A 5-STAR rating is considered excellent. The measures are divided into the following categories:

- Staying Healthy- Includes how often members received various screening tests, vaccines, and other assessments to help them stay healthy.
- Managing chronic conditions- Includes how often members with different long-term conditions completed certain tests and treatments that help manage their conditions; appropriateness of medication and medication adherence.
- Member Satisfaction- Includes ratings of member satisfaction with the plan and how they feel the quality of their care, the plan and providers.
- Customer Service- Includes how responsive and helpful the plan's customer care is and if the information provided is accurate; includes grievances and appeals.
- Plan Operations- Includes how the plan makes decisions about determinations and appeals; formulary pricing; prescription dug safety and findings from CMS audits.

A Plan's STAR ratings are ranked 1-5 in each category, and then used to determine a Plan's overall rating.

\*\*\*\*\*
Above Average Performance
\*\*\*
Average Performance
Below Average Performance
Poor Performance

The STAR rating measures the Plan's performance in over 30 measures covering:

- Clinical quality standards
- Member satisfaction
- Health plan administrative performance
- Compliance with CMS operational standards

Data sources used to measure the performance include:

- Clinical quality standards:
  - Health Effectiveness Data and Information Set (HEDIS®)(HEDIS STARS Measures Link (please see below)
  - Pharmacy Data (Prescription Drug Event-PDE), includes medication adherence
  - Laboratory data
  - Health Outcome Survey (HOS)
     <a href="http://www.hosonline.org/Content/Default.aspx">http://www.hosonline.org/Content/Default.aspx</a>
- Member experience, satisfaction and complaints with Plan:
  - Consumer Assessment of Healthcare Providers and Systems (CAHPS®)
     <a href="https://cahps.ahrq.gov/about-cahps/index.html">https://cahps.ahrq.gov/about-cahps/index.html</a>
  - Medicare Advantage (MA) and Prescription Drug Plan (PBP) CAHPS link <u>http://cms.gov/Research-Statistics-Data-and-</u>
     Systems/Research/CAHPS/mcahps.html
  - Complaint Tracking Module
  - Grievances and Appeals
- Administrative performance and compliance standards
  - CMS Audits
  - Pharmacy (Part D) drug safety and drug pricing accuracy
  - Customer Care Performance

### What is GlobalHealth doing to enhance quality and improve Star ratings?

GlobalHealth's goal is to support our members in maintaining and improving their physical and mental health; effectively manage long-term conditions; and provide high quality, cost effective health care.

We utilize Case and Disease Management, Care Coordination, Quality Improvement and our network providers to encourage our members' healthy choices by monitoring how frequently members receive screenings, vaccines, and other preventive services. Please see GlobalLink<sup>TM</sup> for referral forms.

We have a team dedicated to quality and improving our Star ratings, with a goal of improving the health of our members; attracting new members to our high quality plans and offering competitive reimbursement to our providers. GlobalHealth constantly assesses the Star ratings and the measures that they encompass.

GlobalHealth is also working with specific provider groups in Shared Savings (SSP) agreements, where providers are rewarded for their performance on a number of quality measures, including HEDIS and Stars.

What you can do to continue to provide high quality, timely, appropriate service to our members:

- ✓ Ensure your patients receive routine screenings and preventative services, including labs;
- ✓ Help your patients manage their long- term conditions, including high blood pressure, diabetes, high cholesterol and arthritis;
- ✓ Refer to GlobalHealth Case and Disease Management and Behavioral Health Care Coordination as indicated;
- ✓ Help members choose safe medications (a link to the High Risk Medications list is provided below);
- ✓ Ensure patients are taking their medications as prescribed (principally oral diabetic, cholesterol, HTN, ACE/ ARB);
- ✓ Submit claims and document all services timely, thoroughly and accurately;
- ✓ Understand the impact that <u>you and your staff</u> have on your patient's (our member's) health and satisfaction with their health plan and care, which is reflected in CAHPS and HOS surveys.

# GlobalHealth and Generations Medicare Advantage Plans HEDIS/CAHPS STAR Measures (2015)

### **Bolded** items are GlobalHealth & Generations focus for 2015 Stars

Measure	Description of Measure	4 Star Benchmark
Colorectal Cancer Screening	Medicare members age 50-75 should have a 1) Fecal Occult Blood (gFOBT or iFOBT) within the last year or: 2) flexible sigmoidoscopy within the last 5years or; 3) colonoscopy within the last 10 years. Documentation must include the date test was completed.	≥58%
Breast Cancer Screening	Female Medicare members ages 50- 74 should have a mammography every 2 years.	<u>&gt;</u> 74%
Cholesterol Screening for Patients with Heart Disease	Medicare members ages 18-75 (male and female) who were discharged after an Acute Myocardial Infarction, Coronary Artery Bypass Graft (CABG) or Percutaneous Coronary Interventions (PCI) within the past year, or who have been diagnosed with Ischemic Vascular Disease (IVD) within the past year, should have an LDL-C screening and an LDL-C control (<100 mg/dL) annually.	<u>&gt;</u> 85%
Diabetes Care- Cholesterol Screening	Medicare members ages 18-75 (male and female) with a diagnosis of diabetes with a Cholesterol Screening Test LDL-C, annually.	<u>≥</u> 85%
Annual Flu Vaccine	Medicare members (male and female) who had a flu shot annually prior to flu season.	<u>&gt;</u> 75%
Improving or Maintaining Physical Health	Medicare members (male and female) who state their physical health was the same or better than expected after two years.	<u>≥</u> 60%
Improving or Maintaining Mental Health	Medicare members (male and female) who state their mental health was the same or better than expected after two years.	<u>&gt;</u> 85%
Monitoring Physical Activity	Medicare members age 65 and older (male and female) who discuss exercise with their doctor and were advised to start, increase or maintain their physical activity during the year.	≥60%
Adult BMI Assessment	Medicare members ages 18-74 (male and female) should visit with their primary physician and have a body mass index (BMI) documented every 2 years. Calculate from the member's height and weight.	<u>&gt;</u> 87%
Osteoporosis Management in Women who had a Fracture	Female Medicare members age 67-85 years of age who suffered fracture and received bone mineral density (BMD) test or were prescribed a drug to treat or prevent osteoporosis within 6 months of the fracture. Excludes fractures of fingers, toes, face, and skull. (Refer to Drug list link below)	<u>&gt;</u> 60%

Dishatas Dating I F . F	Madiana manhara assa 10 75 /	>C40/
Diabetes- Retinal Eye Exam	Medicare members ages 18-75 (male and	<u>&gt;</u> 64%
	female) with diabetes (Type 1 and Type 2) who	
	had a dilated or retinal eye exam by an eye care	
	professional to check for damage from diabetes	
	during the year. Results need to indicate date	
Dishetes Kidney Disease	and result.	> OF 0/
Diabetes- Kidney Disease	Medicare members ages 18-75 (male and	<u>&gt;</u> 85%
Monitoring	female) with diabetes who had nephropathy	
	screening test performed every year. a)  Macroalbumin – Any of the following tests done	
	in measurement year: Urinalysis positive	
	(random, spot or timed) for protein, Positive	
	urine dipstick, Positive tablet reagent for urine	
	protein, Positive result for albuminuria, Positive	
	for macroalbuminuria, Positive for proteinuria,	
	Positive for gross proteinuria <i>Note</i> : "trace"	
	urine macroalbumin test results are not	
	considered positive. b) Microalbuminuria –Any	
	of the following tests in measurement year: 24-	
	hour urine for microalbumin Timed urine for	
	microalbumin, Spot urine for microalbumin,	
	Urine for microalbumin/ creatine ratio, 24-hour	
	urine for total protein, Random urine for	
	protein/creatinine ratio or c) has	
	documentation of nephropathy in the record	
	during the past year.	
Diabetes Care- Blood Sugar	Medicare members ages 18-75 (male and	≥80%
Controlled	female) with diabetes who had an HBA1c test in	_
	the measurement year documented in the	
	record with date and results indicating <8.0%.	
Diabetes Care- Cholesterol	Medicare members ages 18-75 (male and	<u>&gt;</u> 64%
Controlled	female) with diabetes who had a cholesterol	
	test in the in the measurement year	
	documented with a LDL-C < 100 mg/ dL)	
Controlling High Blood	Medicare members (male and female)	<u>&gt;</u> 63%
Controlling High Blood Pressure	Medicare members (male and female) diagnosed with HTN, ages 18-59 and all	<u>&gt;</u> 63%
	,	<u>≥</u> 63%
<b>5 5</b>	diagnosed with HTN, ages 18-59 and all members diagnosed with diabetes, with BP of <140-90 mmHg during the measurement year.	≥63%
<b>5 5</b>	diagnosed with HTN, ages 18-59 and all members diagnosed with diabetes, with BP of <140-90 mmHg during the measurement year. Medicare members (male and female)	<u>&gt;</u> 63%
<b>5 5</b>	diagnosed with HTN, ages 18-59 and all members diagnosed with diabetes, with BP of <140-90 mmHg during the measurement year. Medicare members (male and female) diagnosed with HTN, ages 60 and older; with BP	<u>&gt;</u> 63%
<b>5 5</b>	diagnosed with HTN, ages 18-59 and all members diagnosed with diabetes, with BP of <140-90 mmHg during the measurement year. Medicare members (male and female) diagnosed with HTN, ages 60 and older; with BP of <150/90 mmHg during the measurement	<u>&gt;</u> 63%
Pressure	diagnosed with HTN, ages 18-59 and all members diagnosed with diabetes, with BP of <140-90 mmHg during the measurement year. Medicare members (male and female) diagnosed with HTN, ages 60 and older; with BP of <150/90 mmHg during the measurement year.	
Pressure  Rheumatoid Arthritis	diagnosed with HTN, ages 18-59 and all members diagnosed with diabetes, with BP of <140-90 mmHg during the measurement year. Medicare members (male and female) diagnosed with HTN, ages 60 and older; with BP of <150/90 mmHg during the measurement year.  Medicare members (male and female) who	≥63% ≥78%
Pressure	diagnosed with HTN, ages 18-59 and all members diagnosed with diabetes, with BP of <140-90 mmHg during the measurement year. Medicare members (male and female) diagnosed with HTN, ages 60 and older; with BP of <150/90 mmHg during the measurement year.  Medicare members (male and female) who were diagnosed with rheumatoid arthritis	
Pressure  Rheumatoid Arthritis	diagnosed with HTN, ages 18-59 and all members diagnosed with diabetes, with BP of <140-90 mmHg during the measurement year. Medicare members (male and female) diagnosed with HTN, ages 60 and older; with BP of <150/90 mmHg during the measurement year.  Medicare members (male and female) who were diagnosed with rheumatoid arthritis during the measurement year and were	
Pressure  Rheumatoid Arthritis	diagnosed with HTN, ages 18-59 and all members diagnosed with diabetes, with BP of <140-90 mmHg during the measurement year. Medicare members (male and female) diagnosed with HTN, ages 60 and older; with BP of <150/90 mmHg during the measurement year.  Medicare members (male and female) who were diagnosed with rheumatoid arthritis during the measurement year and were dispensed at least one prescription for disease	
Rheumatoid Arthritis Management	diagnosed with HTN, ages 18-59 and all members diagnosed with diabetes, with BP of <140-90 mmHg during the measurement year. Medicare members (male and female) diagnosed with HTN, ages 60 and older; with BP of <150/90 mmHg during the measurement year.  Medicare members (male and female) who were diagnosed with rheumatoid arthritis during the measurement year and were dispensed at least one prescription for disease modifying anti- rheumatic drug (DMARD) filled.	≥78%
Pressure  Rheumatoid Arthritis	diagnosed with HTN, ages 18-59 and all members diagnosed with diabetes, with BP of <140-90 mmHg during the measurement year. Medicare members (male and female) diagnosed with HTN, ages 60 and older; with BP of <150/90 mmHg during the measurement year.  Medicare members (male and female) who were diagnosed with rheumatoid arthritis during the measurement year and were dispensed at least one prescription for disease modifying anti- rheumatic drug (DMARD) filled.  Medicare members 65 years and older (male	
Rheumatoid Arthritis Management	diagnosed with HTN, ages 18-59 and all members diagnosed with diabetes, with BP of <140-90 mmHg during the measurement year. Medicare members (male and female) diagnosed with HTN, ages 60 and older; with BP of <150/90 mmHg during the measurement year.  Medicare members (male and female) who were diagnosed with rheumatoid arthritis during the measurement year and were dispensed at least one prescription for disease modifying anti- rheumatic drug (DMARD) filled.  Medicare members 65 years and older (male and female) who reported having a urine	≥78%
Rheumatoid Arthritis Management	diagnosed with HTN, ages 18-59 and all members diagnosed with diabetes, with BP of <140-90 mmHg during the measurement year. Medicare members (male and female) diagnosed with HTN, ages 60 and older; with BP of <150/90 mmHg during the measurement year.  Medicare members (male and female) who were diagnosed with rheumatoid arthritis during the measurement year and were dispensed at least one prescription for disease modifying anti- rheumatic drug (DMARD) filled.  Medicare members 65 years and older (male and female) who reported having a urine leakage problem in the last six months and who	≥78%
Rheumatoid Arthritis Management	diagnosed with HTN, ages 18-59 and all members diagnosed with diabetes, with BP of <140-90 mmHg during the measurement year. Medicare members (male and female) diagnosed with HTN, ages 60 and older; with BP of <150/90 mmHg during the measurement year.  Medicare members (male and female) who were diagnosed with rheumatoid arthritis during the measurement year and were dispensed at least one prescription for disease modifying anti- rheumatic drug (DMARD) filled.  Medicare members 65 years and older (male and female) who reported having a urine	≥78%

Reducing the Risk of Falling  Plan All-Cause Readmissions	Medicare members 65 years and older (male and female) who had a fall, problems with balance or walking in the past 12 months and who received fall risk intervention from current practitioner. Review of falls and balance annually. Interventions may include recommend use of cane or walker; check BP lying and standing; recommend exercise or physical therapy program; or suggest vision or hearing testing.  Medicare members discharged from hospital	≥59% <9%
	stays who were readmitted to a hospital within 30 days of discharge, either for the same condition as the recent admission <i>or</i> for a different reason.	
Ease of Getting Needed Care and Seeing Specialists	Medicare member report of ease of access to providers, tests and treatment when needed, including care from specialists.	<u>&gt;</u> 85%
Getting Appointments and Care Quickly	Medicare member report of ease of access to providers when needed/ wanted; received care as soon as they thought it was needed and seen within 15 minutes of appointment time.	<u>&gt;</u> 75%
High Risk Medication	Medicare members age 65 and older, who filled 2 or more prescriptions for certain medications with a high risk of serious side effects when safer meds are available. Refer to High Risk Medication List-link below.	<9%
Diabetes Treatment	Medicare members ages 18 and older with HTN and Diabetes who are prescribed and filled at least one prescription for the recommended BP med (ACE inhibitor, ARB DRI therapy) during measurement year.	<u>&gt;</u> 86%
Medication Adherence- Oral Diabetes Medications	Medicare members with diabetes and prescribed an oral diabetic medication, who filled that prescription often enough to cover greater than or equal to 80% of the time they are supposed to be taking the medication.	≥77%
Medication Adherence- Hypertension	Medicare members with HTN and treated with an ACE/ ARB or DRI medication, who filled that prescription often enough to cover greater than or equal to 80% of the time they are supposed to be taking the medication.	≥81%
Medication Adherence- Cholesterol (Statins)	Medicare members treated with a statin cholesterol mediations, who filled that prescription often enough to cover greater than or equal to 80% of the time they are supposed to be taking the medication.	<u>&gt;</u> 76%
Statin Use in Diabetes (new)	Medicare members (male and female) ages 18-75 years of age, diagnosed with HTN and diabetes, who filled at least 2 prescriptions for diabetic medications and at least one prescription for a statin medication during the measurement period.	<u>&gt;</u> 76%

#### **High Risk Medication List:**

http://pqaalliance.org/images/uploads/files/HRM%20Measure%202013website.pdf

Note: The National Committee for Quality Assurance (NCQA) has a comprehensive list of medications and National Drug Code (NDC) codes posted at:

http://www.ncqa.org/HEDISQualityMeasurement/HEDISMeasures/HEDIS2015/HEDIS2015NDCLicense/HEDIS2015FinalNDCLists.aspx