

Prior Authorization Criteria for Drugs on the 2019 State of Oklahoma Retirees (HMO) Formulary

The following are prior authorization criteria for drugs on the 2019 State of Oklahoma Retirees (HMO) formulary. To determine if your drug has a prior authorization requirement, use the GlobalHealth online prescription search specific to your plan at https://globalhealth.com/pharmacy/drug-formularies/. For information on how to request an exception (also called a coverage determination) to the prior authorization criteria for any of these drugs click here.

Updated 09/2019

This document was printed from the GlobalHealth 2019 online formulary. For the most up-to-date listing of covered drugs, call Customer Care at 1-844-280-5555 or, for TTY users, 711, 8:00 a.m. to 8:00 p.m., seven days a week, from October 1 – March 31, and 8:00 a.m. to 8:00 p.m., Monday – Friday, from April 1 – September 30, or visit https://globalhealth.com/pharmacy/drug-formularies/.

PA Criteria

Prior Authorization Group ACITRETIN
Drug Names ACITRETIN

Covered Uses All FDA-approved indications not otherwise excluded from Part D, prevention of

non-melanoma skin cancers in high risk individuals, Lichen planus, Keratosis follicularis

(Darier Disease).

Exclusion Criteria Required Medical Information Age Restrictions Prescriber Restrictions -

Coverage Duration Plan Year

Other Criteria -

Prior Authorization Group ACTIMMUNE
Drug Names ACTIMMUNE

Covered UsesAll FDA-approved indications not otherwise excluded from Part D, mycosis fungoides,

Sezary syndrome, atopic dermatitis.

Exclusion Criteria - Required Medical Information -

Age Restrictions - Prescriber Restrictions -

Coverage Duration Plan Year

Other Criteria -

Prior Authorization GroupADEMPASDrug NamesADEMPAS

Covered Uses All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria

Required Medical Information For pulmonary arterial hypertension (PAH) (WHO Group 1): PAH was confirmed by

right heart catheterization. For chronic thromboembolic pulmonary hypertension (CTEPH) (WHO Group 4): Patient has persistent or recurrent CTEPH after pulmonary endarterectomy (PEA), OR patient has inoperable CTEPH with the diagnosis confirmed by right heart catheterization AND by computed tomography (CT), magnetic resonance

imaging (MRI), or pulmonary angiography. For new starts only (excluding

recurrent/persistent CTEPH after PEA): 1) pretreatment mean pulmonary arterial pressure is greater than or equal to 25 mmHg, AND 2) pretreatment pulmonary capillary wedge pressure is less than or equal to 15 mmHg, AND 3) pretreatment

pulmonary vascular resistance is greater than 3 Wood units.

Age Restrictions -

Prescriber Restrictions -

Coverage Duration Plan Year

Other Criteria -

Drug Names Covered Uses **AFINITOR**

AFINITOR, AFINITOR DISPERZ

All FDA-approved indications not otherwise excluded from Part D, classical Hodgkin

lymphoma, thymomas and thymic carcinomas, Waldenstrom's

macroglobulinemia/lymphoplasmacytic lymphoma, soft tissue sarcoma subtypes:

perivascular epithelioid cell tumors (PEComa), angiomyolipoma,

lymphangioleiomyomatosis, neuroendocrine tumor of the thymus, thyroid carcinoma

(papillary, Hurthle cell, and follicular), osteosarcoma.

Exclusion Criteria

Required Medical Information

For breast cancer: 1) The patient has recurrent or metastatic hormone receptor

positive, HER2 negative disease, 2) Afinitor will be used in combination with

exemestane, and 3) The patient's disease either a) has progressed while on or within 12 months of nonsteroidal aromatase inhibitor therapy, OR b) was previously treated with tamoxifen. For renal cell carcinoma: 1) The disease is relapsed, metastatic or unresectable, and 2) For disease that is of predominantly clear cell histology, disease

has progressed on prior antigiogenic therapy (e.g., sunitinib).

Age Restrictions

Prescriber Restrictions

Coverage Duration

Plan Year

Other Criteria

-

Prior Authorization Group

Drug Names

Covered Uses

Exclusion Criteria

Required Medical Information

AIMOVIC

AIMOVIG

All FDA-approved indications not otherwise excluded from Part D.

1) The patient received at least 3 months of treatment with the requested drug, and the patient had a reduction in migraine days per month from baseline, OR 2) The patient

experienced an inadequate treatment response with a 4-week trial of any of the

following: Antiepileptic drugs (AEDs), Beta-adrenergic blocking agents, Antidepressants. OR 3) The patient experienced an intolerance or has a

contraindication that would prohibit a 4-week trial of any of the following: Antiepileptic

drugs (AEDs), Beta-adrenergic blocking agents, Antidepressants

Age Restrictions

Prescriber Restrictions

Coverage Duration

Initial 3 Months, Reauthorization Plan year

Other Criteria

Prior Authorization GroupALDURAZYMEDrug NamesALDURAZYME

Covered Uses All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria -

Required Medical Information Diagnosis of mucopolysaccharidosis I was confirmed by an enzyme assay

demonstrating a deficiency of alpha-L-iduronidase enzyme activity or by genetic testing. Patients with Scheie syndrome must have moderate to severe symptoms.

Age Restrictions -Prescriber Restrictions --

Coverage Duration Plan Year

Other Criteria -

Prior Authorization GroupALECENSADrug NamesALECENSA

Covered Uses All FDA-approved indications not otherwise excluded from Part D, anaplastic lymphoma

kinase (ALK)-positive recurrent non-small cell lung cancer.

Exclusion Criteria -

Required Medical Information -

Age Restrictions -

Prescriber Restrictions -

Coverage Duration Plan Year

Other Criteria -

Prior Authorization Group ALOSETRON

Drug Names ALOSETRON HYDROCHLORIDE

Covered Uses All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria -

Required Medical Information 1) The requested drug is being prescribed for a biological female or a person that

self-identifies as a female with a diagnosis of severe diarrhea-predominant irritable bowel syndrome (IBS) AND 2) Chronic IBS symptoms lasting at least 6 months AND 3) Gastrointestinal tract abnormalities have been ruled out AND 4) Inadequate response

to conventional therapy.

Age Restrictions - Prescriber Restrictions -

Coverage Duration Plan Year

Other Criteria -

Prior Authorization Group ALPHA1-PROTEINASE INHIBITOR ARALAST NP, PROLASTIN-C, ZEMAIRA **Drug Names**

Covered Uses All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria

Required Medical Information Patients must have clinically evident emphysema. Patients must have a pretreatment

> serum alpha1-proteinase inhibitor level less than 11 micromol/L (80 mg/dl by radial immunodiffusion or 50 mg/dl by nephelometry). Patients must have a pretreatment post-bronchodilation forced expiratory volume in 1 second (FEV1) greater than or equal

to 25 percent and less than or equal to 80 percent of predicted.

Age Restrictions **Prescriber Restrictions**

Coverage Duration Plan Year

Other Criteria

Prior Authorization Group ALUNBRIG ALUNBRIG Drug Names

Covered Uses All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria Required Medical Information Age Restrictions **Prescriber Restrictions**

Coverage Duration Plan Year

Other Criteria

AMPYRA Prior Authorization Group

Drug Names DALFAMPRIDINE ER

Covered Uses All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria

Required Medical Information For new starts: Prior to initiating therapy, patient demonstrates sustained walking

> impairment. For continuation of therapy: Patient must have experienced an improvement in walking speed or other objective measure of walking ability since

starting the requested medication.

Age Restrictions **Prescriber Restrictions**

Coverage Duration Plan Year

Other Criteria

Prior Authorization Group ANADROL
Drug Names ANADROL-50

Covered Uses All FDA-approved indications not otherwise excluded from Part D, Cachexia associated

with AIDS (HIV-wasting), Fanconi's anemia.

Exclusion Criteria -

Required Medical Information

Age Restrictions

Prescriber Restrictions -

Coverage Duration 6 Months

Other Criteria -

Prior Authorization Group APOKYN
Drug Names APOKYN

Covered Uses All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria Required Medical Information Age Restrictions Prescriber Restrictions -

Coverage Duration Plan Year

Other Criteria -

Prior Authorization GroupARCALYSTDrug NamesARCALYST

Covered Uses All FDA-approved indications not otherwise excluded from Part D, prevention of gout

flares in patients initiating or continuing urate-lowering therapy.

Exclusion Criteria -

Required Medical Information For prevention of gout flares in members initiating or continuing urate-lowering therapy

(i.e., allopurinol or febuxostat) (new starts): 1) two or more gout flares within the previous 12 months, AND 2) inadequate response, intolerance or contraindication to maximum tolerated doses of non-steroidal anti-inflammatory drugs and colchicine, AND 3) concurrent use with urate-lowering therapy (i.e., allopurinol or febuxostat). For

prevention of gout flares in members initiating or continuing urate-lowering therapy (i.e., allopurinol or febuxostat) (continuation): 1) member must have achieved or maintained a clinical benefit (i.e., fewer number of gout attacks or fewer flare days) compared to baseline, AND 2) continued use of urate-lowering therapy concurrently with the

requested drug.

Age Restrictions - Prescriber Restrictions -

Coverage Duration For prevention of gout flares: 4 months. Other: Plan Year

Other Criteria -

Prior Authorization GroupARMODAFINILDrug NamesARMODAFINIL

Covered Uses All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria -

Required Medical Information 1) Diagnosis is narcolepsy confirmed by sleep lab evaluation OR 2) Diagnosis is Shift

Work Disorder (SWD) OR 3) Diagnosis is obstructive sleep apnea (OSA) confirmed by

polysomnography

Age Restrictions - Prescriber Restrictions -

Coverage Duration Plan Year

Other Criteria -

Prior Authorization Group AURYXIA
Drug Names AURYXIA

Covered UsesAll FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria Required Medical Information Age Restrictions Prescriber Restrictions -

Coverage Duration Plan Year

Other Criteria -

Prior Authorization GroupAUSTEDODrug NamesAUSTEDO

Covered Uses All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria Required Medical Information Age Restrictions Prescriber Restrictions -

Coverage Duration Plan Year

Other Criteria -

Prior Authorization Group
Drug Names
Covered Uses

AVASTIN AVASTIN

All FDA-approved indications not otherwise excluded from Part D, breast cancer, central nervous system (CNS) tumor types: adult intracranial and spinal ependymoma and anaplastic gliomas, malignant pleural mesothelioma, ovarian malignant sex cord-stromal tumors, soft tissue sarcoma types: AIDS-related Kaposi sarcoma, angiosarcoma and solitary fibrous tumor/hemangiopericytoma, uterine cancer, endometrial cancer, diabetic macular edema, neovascular (wet) age-related macular degeneration including polypoidal choroidopathy and retinal angiomatous proliferation subtypes, macular edema following retinal vein occlusion, proliferative diabetic retinopathy, choroidal neovascularization, neovascular glaucoma, and retinopathy of prematurity.

Exclusion Criteria

Required Medical Information

Age Restrictions

Prescriber Restrictions

Coverage Duration

Other Criteria

-

-

Plan Year

Coverage under Part D will be denied if coverage is available under Part A or Part B as

the medication is prescribed and dispensed or administered for the individual.

Prior Authorization Group Drug Names

B VS. D

ABELCET, ABRAXANE, ACETYLCYSTEINE, ACYCLOVIR SODIUM, ADRIAMYCIN, ADRUCIL, ALBUTEROL SULFATE, ALIMTA, AMBISOME, AMINOSYN II, AMINOSYN-PF, AMINOSYN-PF 7%, AMPHOTERICIN B, APREPITANT, AZACITIDINE, AZATHIOPRINE, BENDEKA, BLEOMYCIN SULFATE, BUDESONIDE, CALCITONIN-SALMON, CALCITRIOL, CARBOPLATIN, CINACALCET HYDROCHLORIDE, CISPLATIN, CLINIMIX 4.25%/DEXTROSE 1, CLINIMIX 4.25%/DEXTROSE 2. CLINIMIX 4.25%/DEXTROSE 5. CLINIMIX 5%/DEXTROSE 15%. CLINIMIX 5%/DEXTROSE 20%. CLINIMIX 5%/DEXTROSE 25%. CROMOLYN SODIUM, CYCLOPHOSPHAMIDE, CYCLOSPORINE, CYCLOSPORINE MODIFIED. CYTARABINE AQUEOUS, DACARBAZINE, DEPO-PROVERA, DEXRAZOXANE, DIPHTHERIA/TETANUS TOXOID, DOCETAXEL, DOXORUBICIN HCL. DOXORUBICIN HYDROCHLORIDE, DRONABINOL, EMEND, ENGERIX-B. EPIRUBICIN HCL, ETOPOSIDE, FASLODEX, FLUOROURACIL, FREAMINE HBC 6.9%, FREAMINE III, FULVESTRANT, GAMASTAN S/D, GANCICLOVIR, GEMCITABINE, GEMCITABINE HCL, GENGRAF, GRANISETRON HCL, HEPARIN SODIUM, HEPATAMINE, HUMULIN R U-500 (CONCENTR, HYDROMORPHONE HCL, IBANDRONATE SODIUM, IFEX, IFOSFAMIDE, IMOVAX RABIES (H.D.C.V.). INTRALIPID, INTRON A, IPRATROPIUM BROMIDE, IPRATROPIUM BROMIDE/ALBUT, IRINOTECAN, KADCYLA, LEUCOVORIN CALCIUM, LEVALBUTEROL, LEVALBUTEROL HCL, LEVOCARNITINE, LIDOCAINE HCL, METHOTREXATE SODIUM, METHYLPREDNISOLONE, METHYLPREDNISOLONE ACETAT, METHYLPREDNISOLONE SODIUM, MITOMYCIN, MORPHINE SULFATE, MYCOPHENOLATE MOFETIL, MYCOPHENOLIC ACID DR, NEBUPENT, NEPHRAMINE, NULOJIX, NUTRILIPID, ONDANSETRON HCL, ONDANSETRON HYDROCHLORIDE, ONDANSETRON ODT, OXALIPLATIN, PACLITAXEL. PAMIDRONATE DISODIUM, PARICALCITOL, PREDNISOLONE, PREDNISOLONE SODIUM PHOSP, PREDNISONE, PREDNISONE INTENSOL, PREMASOL. PROCALAMINE, PROGRAF, PROSOL, RABAVERT, RAPAMUNE, RECOMBIVAX HB, SANDIMMUNE, SENSIPAR, SIROLIMUS, TACROLIMUS, TAXOTERE, TDVAX, TENIVAC, TOPOSAR, TOPOTECAN HCL, TOPOTECAN HYDROCHLORIDE, TPN ELECTROLYTES, TRAVASOL, TROPHAMINE, VINBLASTINE SULFATE, VINCASAR PFS, VINCRISTINE SULFATE, VINORELBINE TARTRATE, XATMEP, ZOLEDRONIC ACID. ZORTRESS

Covered Uses

This drug may be covered under Medicare Part B or D depending upon the circumstances. Information may need to be submitted describing the use and setting of the drug to make the determination.

Exclusion Criteria Required Medical Information Age Restrictions Prescriber Restrictions Coverage Duration N/A
Other Criteria -

Prior Authorization GroupBALVERSADrug NamesBALVERSA

Covered Uses All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria Required Medical Information Age Restrictions Prescriber Restrictions -

Coverage Duration Plan Year

Other Criteria -

Prior Authorization Group BANZEL Drug Names BANZEL

Covered Uses All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria - Required Medical Information -

Age Restrictions 1 year of age or older.

Prescriber Restrictions -

Coverage Duration Plan Year

Other Criteria -

Prior Authorization GroupBENLYSTADrug NamesBENLYSTA

Covered Uses All FDA-approved indications not otherwise excluded from Part D.

Exclusion CriteriaSevere active lupus nephritis. Severe active central nervous system lupus. **Required Medical Information**Patient has been diagnosed with active, autoantibody-positive systemic lupus

erythematosus (SLE). Patient is currently receiving standard therapy for SLE (e.g., corticosteroids, azathioprine, leflunomide, methotrexate, mycophenolate mofetil, hydroxychloroquine, non-steroidal anti-inflammatory drugs) OR patient is not currently receiving standard therapy for SLE because patient tried and had an inadequate

response or intolerance to standard therapy.

Age Restrictions - Prescriber Restrictions -

Coverage Duration Plan Year

Other Criteria -

Prior Authorization GroupBERINERTDrug NamesBERINERT

Covered Uses All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria -

Required Medical Information Patient has hereditary angioedema (HAE) with C1 inhibitor deficiency confirmed by

laboratory testing OR patient has hereditary angioedema with normal C1 inhibitor confirmed by laboratory testing. For patients with HAE with normal C1 inhibitor,

EITHER 1) Patient tested positive for the F12 gene mutation OR 2) Patient has a family history of angioedema and the angioedema was refractory to a trial of antihistamine for

at least one month.

Age Restrictions -

Prescriber Restrictions

Coverage Duration Plan Year

Other Criteria -

Prior Authorization GroupBETASERONDrug NamesBETASERON

Covered Uses All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria -

Required Medical Information Have a relapsing form of multiple sclerosis (MS) (e.g., relapsing-remitting MS,

progressive-relapsing MS, or secondary progressive MS with relapses) OR first clinical

episode of MS.

Age Restrictions -

Prescriber Restrictions -

Coverage Duration Plan Year

Other Criteria -

Prior Authorization Group BEXAROTENE

Drug Names BEXAROTENE, TARGRETIN

Covered Uses All FDA-approved indications not otherwise excluded from Part D, mycosis fungoides,

Sezary syndrome (capsules only), primary cutaneous CD30-positive T-cell

lymphoproliferative disorder types: primary cutaneous anaplastic large cell lymphoma (capsules only) and lymphomatoid papulosis (capsules only), chronic or smoldering adult T-cell leukemia/lymphoma (gel only), primary cutaneous B-cell lymphoma types: primary cutaneous marginal zone lymphoma (gel only) and primary cutaneous follicle

center lymphoma (gel only).

Exclusion Criteria -

Required Medical Information - Age Restrictions -

Prescriber Restrictions

Coverage Duration Plan Year

Other Criteria -

Prior Authorization Group BOSENTAN

Drug Names BOSENTAN, TRACLEER

Covered Uses All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria -

Required Medical Information Pulmonary arterial hypertension (WHO Group 1) was confirmed by right heart

catheterization. For new starts only: 1) pretreatment mean pulmonary arterial pressure is greater than or equal to 25 mmHg, 2) pretreatment pulmonary capillary wedge pressure is less than or equal to 15 mmHg, and 3) pretreatment pulmonary vascular

resistance is greater than 3 Wood units.

Age Restrictions -

Prescriber Restrictions -

Coverage Duration Plan Year

Other Criteria -

Prior Authorization Group BOSULIF

Drug Names BOSULIF

Covered Uses All FDA-approved indications not otherwise excluded from Part D, relapsed/refractory

Philadelphia chromosome positive acute lymphoblastic leukemia (Ph+ ALL).

Exclusion Criteria

Required Medical Information Diagnosis of CML was confirmed by detection of the Philadelphia chromosome or

BCR-ABL gene. For CML, patient meets one of the following: 1) Patient has chronic phase CML, OR 2) Patient has accelerated or blast phase CML, OR 3) Patient received

a hematopoietic stem cell transplant.

Age Restrictions -

Prescriber Restrictions -

Coverage Duration Plan Year

Other Criteria -

Prior Authorization Group BRAFTOVI Drug Names BRAFTOVI

Covered Uses All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria -

Required Medical Information -

Age Restrictions -

Prescriber Restrictions

Coverage Duration Plan Year

Other Criteria -

Prior Authorization GroupBRIVIACTDrug NamesBRIVIACT

Covered Uses All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria - Required Medical Information -

Age Restrictions 4 years of age or older (tablets and oral solution).

Prescriber Restrictions -

Coverage Duration Plan Year

Other Criteria -

Prior Authorization GroupBUPRENORPHINEDrug NamesBUPRENORPHINE HCL

Covered Uses All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria -

Required Medical Information 1) The requested drug is being prescribed for the treatment of opioid dependence AND

2) If the patient is pregnant or breastfeeding and being prescribed the requested drug for induction therapy and/or subsequent maintenance therapy for opioid dependence treatment OR 3) If the requested drug is being prescribed for induction therapy for transition from opioid use to opioid dependence treatment OR 4) If the requested drug is being prescribed for maintenance therapy for opioid dependence treatment in a

patient who is intolerant to naloxone.

Coverage Duration 12 Months

Other Criteria -

Prior Authorization GroupCABOMETYXDrug NamesCABOMETYX

Covered Uses All FDA-approved indications not otherwise excluded from Part D, non-small cell lung

cancer.

Exclusion Criteria -

Required Medical Information For renal cell carcinoma: The disease is relapsed, unresectable, or metastatic.

Age Restrictions Prescriber Restrictions -

Coverage Duration Plan Year

Other Criteria -

Prior Authorization Group CALCIPOTRIENE

Drug Names CALCIPOTRIENE, CALCITRENE, ENSTILAR

Covered Uses All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria -

Required Medical Information 1) The requested drug is being prescribed for the treatment of psoriasis AND 2) The

patient experienced an inadequate treatment response, intolerance, or contraindication

to a generic topical steroid.

Age Restrictions -Prescriber Restrictions --

Coverage Duration Plan Year

Other Criteria -

Prior Authorization Group CALQUENCE
Drug Names CALQUENCE

Covered Uses All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria Required Medical Information Age Restrictions Prescriber Restrictions -

Coverage Duration Plan Year

Other Criteria -

Prior Authorization Group CAPRELSA Drug Names CAPRELSA

Covered UsesAll FDA-approved indications not otherwise excluded from Part D, non-small lung

cancer and differentiated thyroid carcinoma: papillary, follicular, Hurthle cell.

Exclusion Criteria -

Required Medical Information For non-small cell lung cancer (NSCLC): the requested drug is used for NSCLC with

RET gene rearrangements.

Age Restrictions -

Prescriber Restrictions -

Coverage Duration Plan Year

Other Criteria -

Prior Authorization Group CARBAGLU
Drug Names CARBAGLU

Covered UsesAll FDA-approved indications not otherwise excluded from Part D, methylmalonic

acidemia, propionic acidemia.

Exclusion Criteria

Required Medical Information Diagnosis of NAGS deficiency was confirmed by enzymatic or genetic testing.

Age Restrictions -

Prescriber Restrictions

Coverage Duration Plan Year

Other Criteria -

Prior Authorization GroupCAYSTONDrug NamesCAYSTON

Covered Uses All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria -

Required Medical Information Pseudomonas aeruginosa is present in the patient's airway cultures OR the patient has

a history of pseudomonas aeruginosa infection or colonization in the airways.

Age Restrictions - Prescriber Restrictions -

Coverage Duration Plan Year

Other Criteria -

Prior Authorization Group CERDELGA
Drug Names CERDELGA

Covered Uses All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria -

Required Medical Information Diagnosis of Gaucher disease was confirmed by an enzyme assay demonstrating a

deficiency of beta-glucocerebrosidase enzyme activity or by genetic testing. The patient's CYP2D6 metabolizer status has been established using an FDA-cleared test. The patient is a CYP2D6 extensive metabolizer, an intermediate metabolizer, or a poor

metabolizer.

Age Restrictions 18 years of age or older

Prescriber Restrictions

Coverage Duration Plan Year

Other Criteria -

Prior Authorization Group CEREZYME
Drug Names CEREZYME

Covered UsesAll FDA-approved indications not otherwise excluded from Part D, type 3 Gaucher

disease.

Exclusion Criteria

Required Medical Information Diagnosis of Gaucher disease was confirmed by an enzyme assay demonstrating a

deficiency of beta-glucocerebrosidase enzyme activity or by genetic testing.

Age Restrictions -

Prescriber Restrictions -

Coverage Duration Plan Year

Other Criteria -

Prior Authorization Group CHANTIX

Drug Names CHANTIX, CHANTIX CONTINUING MONTH, CHANTIX STARTING MONTH PA

Covered Uses All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria Required Medical Information Age Restrictions Prescriber Restrictions -

Coverage Duration 6 Months

Other Criteria -

Prior Authorization Group CLOMIPRAMINE
Drug Names CLOMIPRAMINE HCL

Covered UsesAll FDA-approved indications not otherwise excluded from Part D, Depression, Panic

Disorder.

Exclusion Criteria -

Required Medical Information 1) Obsessive-Compulsive Disorder (OCD) or Panic Disorder AND 2) The patient has

of the following: a generic selective serotonin reuptake inhibitor (SSRI), a generic serotonin and norepinephrine reuptake inhibitor (SNRI), mirtazapine OR 3) Depression AND 4) The patient has experienced an inadequate treatment response, intolerance or

experienced an inadequate treatment response, intolerance or contraindication to one

contraindication to one of the following: a generic selective serotonin reuptake inhibitor (SSRI), a generic serotonin and norepinephrine reuptake inhibitor (SNRI), mirtazapine,

bupropion.

Coverage Duration Plan Year

Other Criteria -

Prior Authorization Group CLORAZEPATE

Drug Names CLORAZEPATE DIPOTASSIUM

Covered Uses All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria -

Required Medical Information 1) For the management of anxiety disorders, the requested drug is being used with a

selective serotonin reuptake inhibitor (SSRI) or serotonin-norepinephrine reuptake inhibitor (SNRI) until the antidepressant becomes effective for the symptoms of anxiety OR the patient has experienced an inadequate treatment response, intolerance or

contraindication to a selective serotonin reuptake inhibitor (SSRI) or a

serotonin-norepinephrine reuptake inhibitor (SNRI) OR 2) For adjunctive therapy in the management of partial seizures OR 3) Symptomatic relief in acute alcohol withdrawal OR 4) For the short-term relief of the symptoms of anxiety AND 5) The benefit of therapy with the prescribed medication outweighs the potential risk in a patient 65

years of age or older.

Age Restrictions
Prescriber Restrictions

Coverage Duration Short-term relief anxiety-1 Month, Anxiety Disorders-4 Months, All other

Diagnoses-Plan Year

Other Criteria This Prior Authorization requirement only applies to patients 65 years of age or older.

The American Geriatrics Society identifies the use of this medication as potentially inappropriate in older adults, meaning it is best avoided, prescribed at reduced dosage,

or used with caution or carefully monitored.

Prior Authorization Group CLOZAPINE ODT CLOZAPINE ODT CLOZAPINE ODT

Covered Uses All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria Required Medical Information Age Restrictions Prescriber Restrictions -

Coverage Duration Plan Year

Other Criteria -

Prior Authorization Group COMETRIQ
Drug Names COMETRIQ

Covered Uses All FDA-approved indications not otherwise excluded from Part D, non-small lung

cancer and differentiated thyroid carcinoma: papillary, follicular, Hurthle cell.

Exclusion Criteria

Required Medical Information For non-small cell lung cancer (NSCLC): The requested drug is used for NSCLC with

RET gene rearrangements.

Age Restrictions -Prescriber Restrictions --

Coverage Duration Plan Year

Other Criteria -

Prior Authorization GroupCOPIKTRADrug NamesCOPIKTRA

Covered Uses All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria Required Medical Information Age Restrictions Prescriber Restrictions -

Coverage Duration Plan Year

Other Criteria -

Prior Authorization Group COTELLIC Drug Names COTELLIC

Covered Uses All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria Required Medical Information Age Restrictions Prescriber Restrictions -

Coverage Duration Plan Year

Other Criteria -

Prior Authorization Group CYSTAGON Drug Names CYSTAGON

Covered Uses All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria -

Required Medical Information Diagnosis of nephropathic cystinosis was confirmed by the presence of increased

cystine concentration in leukocytes or by genetic testing.

Age Restrictions - Prescriber Restrictions -

Coverage Duration Plan Year

Other Criteria -

Prior Authorization Group CYSTARAN
Drug Names CYSTARAN

Covered Uses All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria -

Required Medical Information Diagnosis of cystinosis was confirmed by the presence of increased cystine

concentration in leukocytes or by DNA testing. The patient has corneal cystine crystal

accumulation.

Age Restrictions - Prescriber Restrictions -

Coverage Duration Plan Year

Other Criteria -

Prior Authorization GroupDAURISMODrug NamesDAURISMO

Covered Uses All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria Required Medical Information Age Restrictions Prescriber Restrictions -

Coverage Duration Plan Year

Other Criteria -

Prior Authorization Group DEFERASIROX

Drug Names JADENU, JADENU SPRINKLE

Covered Uses All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria -

Required Medical Information For chronic iron overload due to blood transfusions: pretreatment serum ferritin level is

greater than 1000 mcg/L.

Age Restrictions --

Coverage Duration Plan Year

Other Criteria -

Prior Authorization GroupDEMSERDrug NamesDEMSER

Covered Uses All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria Required Medical Information Age Restrictions Prescriber Restrictions -

Coverage Duration Plan Year

Other Criteria -

Prior Authorization GroupDESVENLAFAXINEDrug NamesDESVENLAFAXINE ER

Covered Uses All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria -

Required Medical Information Patient experienced an inadequate treatment response, intolerance, or contraindication

to any of the following: a generic serotonin and norepinephrine reuptake inhibitor (SNRI), a generic selective serotonin reuptake inhibitor (SSRI), mirtazapine, bupropion.

Age Restrictions -

Prescriber Restrictions -

Coverage Duration Plan Year

Other Criteria -

Prior Authorization Group DIAZEPAM

Drug Names DIAZEPAM, DIAZEPAM INTENSOL

Covered Uses All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria -

Required Medical Information 1) For the management of anxiety disorders, the requested drug is being used with a

selective serotonin reuptake inhibitor (SSRI) or serotonin-norepinephrine reuptake inhibitor (SNRI) until the antidepressant becomes effective for the symptoms of anxiety OR the patient has experienced an inadequate treatment response, intolerance or

contraindication to a selective serotonin reuptake inhibitor (SSRI) or a

serotonin-norepinephrine reuptake inhibitor (SNRI) OR 2) For symptomatic relief in acute alcohol withdrawal OR 3) For use as an adjunct for the relief of skeletal muscle spasms OR 4) For adjunctive therapy in the treatment of convulsive disorders OR 5) For the short-term relief of the symptoms of anxiety AND 6) The benefit of therapy with the prescribed medication outweighs the potential risk in a patient 65 years of age or

older.

Age Restrictions
Prescriber Restrictions

Coverage Duration

Short-term relief anxiety-1 Month, Anxiety Disorders-4 Months, All other

Diagnoses-Plan Year

Other Criteria This Prior Authorization requirement only applies to patients 65 years of age or older.

The American Geriatrics Society identifies the use of this medication as potentially inappropriate in older adults, meaning it is best avoided, prescribed at reduced dosage,

or used with caution or carefully monitored

Prior Authorization GroupDICLOFENAC GEL 1%Drug NamesDICLOFENAC SODIUM

Covered Uses All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria -

Required Medical Information 1) The patient has osteoarthritis pain in joints susceptible to topical treatment such as

feet, ankles, knees, hands, wrist, or elbow AND 2) Treatment with the requested drug is

necessary due to intolerance or a contraindication to oral nonsteroidal

anti-inflammatory (NSAID) drugs.

Age Restrictions -

Prescriber Restrictions -

Coverage Duration Plan Year

Other Criteria -

Prior Authorization Group EMGALITY
Drug Names EMGALITY

Covered Uses All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria -

Required Medical Information 1) The patient received at least 3 months of treatment with the requested drug, and the

patient had a reduction in migraine days per month from baseline, OR 2) The patient experienced an inadequate treatment response with a 4-week trial of any of the

following: Antiepileptic drugs (AEDs), Beta-adrenergic blocking agents, Antidepressants, OR 3) The patient experienced an intolerance or has a

contraindication that would prohibit a 4-week trial of any of the following: Antiepileptic

drugs (AEDs), Beta-adrenergic blocking agents, Antidepressants

Age Restrictions - Prescriber Restrictions -

Coverage Duration Initial 3 Months, Reauthorization Plan year

Other Criteria -

Prior Authorization Group EMSAM
Drug Names EMSAM

Covered Uses All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria -

Required Medical Information 1) Patient experienced an inadequate treatment response, intolerance, or

contraindication to any of the following antidepressants: bupropion, trazodone,

mirtazapine, serotonin norepinephrine reuptake inhibitors (e.g., venlafaxine), selective serotonin reuptake inhibitors (e.g., citalopram, fluoxetine, fluvoxamine, paroxetine, sertraline), tricyclic or tetracyclic antidepressants (e.g., amitriptyline, nortriptyline) OR 2)

Patient is unable to swallow oral formulations.

Age Restrictions 18 years of age or older.

Prescriber Restrictions -

Coverage Duration Plan Year

Other Criteria -

Prior Authorization Group ENDARI Drug Names ENDARI

Covered Uses All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria - Required Medical Information -

Age Restrictions 5 years of age or older

Prescriber Restrictions -

Coverage Duration Plan Year

Other Criteria -

Prior Authorization GroupEPCLUSADrug NamesEPCLUSA

Covered Uses All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria -

Required Medical Information Chronic hepatitis C infection confirmed by presence of HCV RNA in the serum prior to

starting treatment. Planned treatment regimen, genotype, prior treatment history, presence or absence of cirrhosis (compensated or decompensated [Child Turcotte Pugh class B or C]), presence or absence of HIV coinfection, presence or absence of resistance-associated substitutions where applicable, liver and kidney transplantation status if applicable. Coverage conditions and specific durations of approval will be

based on current AASLD treatment guidelines.

Age Restrictions -

Prescriber Restrictions -

Coverage Duration Criteria will be applied consistent with current AASLD-IDSA guidance.

Other Criteria -

Prior Authorization GroupEPIDIOLEXDrug NamesEPIDIOLEX

Covered Uses All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria Required Medical Information Age Restrictions Prescriber Restrictions -

Coverage Duration Plan Year

Other Criteria -

Prior Authorization Group Drug Names Covered Uses

FPO **PROCRIT**

All FDA-approved indications not otherwise excluded from Part D, anemia due to myelodysplastic syndromes (MDS), anemia in congestive heart failure (CHF), anemia in rheumatoid arthritis (RA), anemia due to hepatitis C treatment (ribavirin in combination with either interferon alfa or peginterferon alfa), anemia in primary myelofibrosis, post-polycythemia vera myelofibrosis, and post-essential thrombocythemia myelofibrosis.

Exclusion Criteria Required Medical Information Patients receiving chemotherapy with curative intent. Patients with myeloid cancer. For all uses except surgery: Pretreatment (no erythropoietin treatment in previous month) Hgb is less than 10 g/dL (less than 9 g/dL for anemia in CHF only). Additional requirements for primary myelofibrosis (MF), post-polycythemia vera MF, post-essential thrombocythemia MF: 1) Patient has symptomatic anemia and 2) For initial therapy, pretreatment serum erythropoietin level is less than 500mU/mL. For surgery: 1) Patient is scheduled for elective, noncardiac, nonvascular surgery and 2) Pretreatment Hgb is greater than 10 but not more than 13 g/dL.

Age Restrictions **Prescriber Restrictions Coverage Duration** Other Criteria

16 weeks

Coverage under Part D will be denied if coverage is available under Part A or Part B as the medication is prescribed and dispensed or administered for the individual (e.g., used for treatment of anemia for a patient with chronic renal failure who is undergoing dialysis, or furnished from physician's supply incident to a physician service). Coverage includes use in anemia in patients whose religious beliefs forbid blood transfusions. Requirements regarding Hgb values exclude values due to a recent transfusion. For reauthorizations (patient received erythropoietin in previous month): 1) For all uses except surgery, there is an increase in Hgb of at least 1 g/dL after at least 12 weeks of therapy, 2) For anemia in CKD, MDS, CHF, RA, HIV, hepatitis C treatment, primary MF, post-polycythemia vera MF, post-essential thrombocythemia MF, or patients whose religious beliefs forbid blood transfusions: current Hgb is less than or equal to 12 g/dL, and 3) For anemia due to myelosuppressive cancer chemotherapy: current Hgb is less than 11 g/dL.

Prior Authorization Group

ERIVEDGE ERIVEDGE Drug Names

Covered Uses

All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria

Required Medical Information Age Restrictions

Prescriber Restrictions

Plan Year

Coverage Duration Other Criteria

Prior Authorization Group ERLEADA
Drug Names ERLEADA

Covered Uses All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria -

Required Medical Information The requested drug will be used in combination with a gonadotropin-releasing hormone

(GnRH) analog or after bilateral orchiectomy.

Age Restrictions -

Prescriber Restrictions -

Coverage Duration Plan Year

Other Criteria -

Prior Authorization GroupESBRIETDrug NamesESBRIET

Covered Uses All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria -

Required Medical Information Initial Review Only: The patient does not have a known etiology for interstitial lung

disease and meets one of the following: 1) a high-resolution computed tomography

(HRCT) study of the chest or surgical lung biopsy reveals the usual interstitial pneumonia (UIP) pattern, or 2) HRCT study of the chest reveals a possible UIP pattern and the diagnosis is supported either by surgical lung biopsy or by a multidisciplinary discussion between at least a radiologist and pulmonologist who are experienced in idiopathic pulmonary fibrosis if surgical lung biopsy has not been conducted. For

continuation: The patient does not have a known etiology for interstitial lung disease.

Age Restrictions -

Prescriber Restrictions -

Coverage Duration Plan Year

Other Criteria -

Prior Authorization GroupFABRAZYMEDrug NamesFABRAZYME

Covered Uses All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria -

Required Medical Information Diagnosis of Fabry disease was confirmed by an enzyme assay demonstrating a

deficiency of alpha-galactosidase enzyme activity or by genetic testing, or the patient is

an obligate female carrier with a first degree male relative diagnosed with Fabry

disease.

Age Restrictions -

Prescriber Restrictions -

Coverage Duration Plan Year

Other Criteria -

Prior Authorization Group FARYDAK **Drug Names** FARYDAK

Covered Uses All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria Required Medical Information Age Restrictions Prescriber Restrictions -

Coverage Duration Plan Year

Other Criteria -

Prior Authorization Group FENTANYL PATCH

Drug Names FENTANYL

Covered Uses All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria -

Required Medical Information 1) The requested drug is being prescribed for pain associated with cancer, sickle cell

disease, a terminal condition, or pain being managed through palliative care OR 2) The

requested drug is being prescribed for pain severe enough to require daily, around-the-clock, long-term treatment in a patient who has been taking an opioid [Note: This drug should be prescribed only by healthcare professionals who are knowledgeable in the use of potent opioids for the management of chronic pain.] AND

3) The patient can safely take the requested dose based on their history of opioid use AND 4) The patient has been evaluated and will be monitored for the development of

opioid use disorder

Age Restrictions -

Prescriber Restrictions -

Coverage Duration Plan Year

Other Criteria -

Prior Authorization Group FETZIMA

Drug Names FETZIMA, FETZIMA TITRATION PACK

Covered Uses All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria -

Required Medical Information Patient experienced an inadequate treatment response, intolerance, or contraindication

to two generic alternatives from the following drug classes: selective serotonin reuptake

inhibitors (SSRIs) or serotonin-norepinephrine reuptake inhibitors (SNRIs).

Age Restrictions -

Prescriber Restrictions -

Coverage Duration Plan Year

Other Criteria -

Prior Authorization Group FIRAZYR **Drug Names** FIRAZYR

Covered Uses All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria -

Required Medical Information The requested drug is being used for the treatment of acute angioedema attacks.

Patient has hereditary angioedema (HAE) with C1 inhibitor deficiency confirmed by laboratory testing OR patient has hereditary angioedema with normal C1 inhibitor confirmed by laboratory testing. For patients with HAE with normal C1 inhibitor,

EITHER 1) Patient tested positive for the F12 gene mutation OR 2) Patient has a family history of angioedema and the angioedema was refractory to a trial of antihistamine for

at least one month.

Age Restrictions 18 years of age or older

Prescriber Restrictions -

Coverage Duration Plan Year

Other Criteria -

Prior Authorization GroupFORTEODrug NamesFORTEO

Covered Uses All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria -

Required Medical Information For postmenopausal osteoporosis: patient has ONE of the following (1. or 2.): 1) A

history of fragility fractures, OR 2) A pre-treatment T-score of less than or equal to -2.5 or osteopenia with a high pre-treatment FRAX fracture probability and patient has ANY of the following: a) Indicators for higher fracture risk (e.g., advanced age, frailty, glucocorticoid therapy, very low T-scores, or increased fall risk), OR b) Patient has failed prior treatment with or is intolerant to a previous osteoporosis therapy (i.e., oral bisphosphonates or injectable antiresorptive agents). For primary or hypogonadal osteoporosis in men: patient has a) a history of osteoporotic vertebral or hip fracture OR b) a pre-treatment T-score of less than or equal to -2.5 OR c) osteopenia with a high pre-treatment FRAX fracture probability. For glucocorticoid-induced osteoporosis: Patient has had an oral bisphosphonate trial of at least 1-year duration unless patient has a contraindication or intolerance to an oral bisphosphonate, AND Patient has a) a history of fragility fracture, OR b) a pre-treatment T-score of less than or equal to -2.5,

OR c) osteopenia with a high pre-treatment FRAX fracture probability.

Age Restrictions Prescriber Restrictions Coverage Duration 24 months (lifetime)

Other Criteria

Patient has high FRAX fracture probability if the 10 year probability is either greater than or equal to 20% for any major osteoporotic fracture or greater than or equal to 3%

for hip fracture

Prior Authorization GroupFYCOMPADrug NamesFYCOMPA

Covered Uses All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria - Required Medical Information -

Age Restrictions Partial-onset seizures - 4 years of age or older. Primary generalized tonic-clonic

seizures - 12 years of age or older.

Prescriber Restrictions

Coverage Duration Plan Year

Other Criteria -

Prior Authorization Group GATTEX **Drug Names** GATTEX

Covered Uses All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria -

Required Medical Information For initial therapy: Patient was dependent on parenteral support for at least 12 months.

For continuation: Requirement for parenteral support has decreased from baseline

while on teduglutide therapy.

Age Restrictions -

Prescriber Restrictions -

Coverage Duration Plan Year

Other Criteria -

Prior Authorization GroupGILENYADrug NamesGILENYA

Covered Uses All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria -

Required Medical Information Have a relapsing form of multiple sclerosis (MS) (e.g., relapsing-remitting MS,

progressive-relapsing MS, or secondary progressive MS with relapses).

Age Restrictions -

Prescriber Restrictions -

Coverage Duration Plan Year

Other Criteria -

Prior Authorization GroupGILOTRIFDrug NamesGILOTRIF

Covered Uses All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria -

Required Medical Information For non-small cell lung cancer (NSCLC), patient meets either of the following: 1) patient

has metastatic squamous NSCLC that progressed after platinum-based chemotherapy,

OR 2) patient has a known sensitizing epidermal growth factor receptor (EGFR)

mutation.

Age Restrictions -

Prescriber Restrictions -

Coverage Duration Plan Year

Other Criteria -

Prior Authorization Group GLATIRAMER

Drug Names GLATIRAMER ACETATE, GLATOPA

Covered Uses All FDA-approved indications not otherwise excluded from Part D, first clinical episode

of MS.

Exclusion Criteria -

Required Medical Information Have a relapsing form of multiple sclerosis (MS) (e.g., relapsing-remitting MS,

progressive-relapsing MS, or secondary progressive MS with relapses) OR first clinical

episode of MS.

Age Restrictions -

Prescriber Restrictions -

Coverage Duration Plan Year

Other Criteria -

Prior Authorization Group GRANIX

Drug Names GRANIX

Covered UsesAll FDA-approved indications not otherwise excluded from Part D, treatment of

chemotherapy-induced febrile neutropenia (FN), stem cell transplantation related indications, acute lymphocytic leukemia (ALL), acute myeloid leukemia (AML), severe chronic neutropenia (congenital, cyclic, or idiopathic), myelodysplastic syndromes (MDS), agranulocytosis, aplastic anemia, HIV-related neutropenia, neutropenia related

to renal transplantation.

Exclusion Criteria Use of the requested product within 24 hours prior to or following chemotherapy.

Required Medical Information For prophylaxis or treatment of myelosuppressive chemotherapy-induced FN patients

must meet all of the following: 1) Patient has a non-myeloid cancer, 2) Patient has received, is currently receiving, or will be receiving treatment with myelosuppressive

anti-cancer therapy

Coverage Duration 6 months

Other Criteria -

Drug Names Covered Uses

Exclusion Criteria

Required Medical Information

GROWTH HORMONE

GENOTROPIN, GENOTROPIN MINIQUICK

All medically accepted indications not otherwise excluded from Part D.

Pediatric nationts with closed eniphyses (except in nationts with PWS)

Pediatric patients with closed epiphyses (except in patients with PWS).

Pediatric GHD: 1) Younger than 2.5 yrs old, when applicable: a) Pre-treatment (pre-tx) height (ht) more than 2 SD below mean and slow growth velocity. 2) 2.5 yrs old or older: a) Pre-tx 1-year ht velocity more than 2 SD below mean OR b) Pre-tx ht more than 2 SD below mean and 1-year ht velocity more than 1 SD below mean. Pediatric GHD: 1) Failed 2 stimulation tests (peak below 10 ng/mL) prior to starting treatment, OR 2) Pituitary/CNS disorder (eg. genetic defects, CNS tumors, congenital structural abnormalities) and pre-tx IGF-1 more than 2 SD below mean. OR 3) Patient is a neonate or was diagnosed with GHD as a neonate. TS: 1) Confirmed by karyotyping AND 2) Pre-treatment height is less than the 5th percentile for age. SGA: 1) Birth weight (wt) below 2500g at gestational age (GA) more than 37 weeks OR birth wt or length below 3rd percentile for GA or at least 2 SD below mean for GA. AND 2) Did not manifest catch-up growth by age 2. Adult GHD: 1) Failed 2 stimulation tests (peak below 5 ng/mL) or test with Macrilen (peak below 2.8 ng/ml) prior to starting tx, OR 2) Structural abnormality of the hypothalamus/pituitary AND 3 or more pituitary hormone deficiencies, OR 3) Childhood-onset GHD with congenital (genetic or structural) abnormality of the hypothalamus/pituitary/CNS, OR 4) Low pre-tx IGF-1 and failed 1 stimulation test prior to starting tx.

Age Restrictions
Prescriber Restrictions

SGA: 2 years of age or older

Endocrinologist, pediatric endocrinologist, pediatric nephrologist, infectious disease

specialist, gastroenterologist/nutritional support specialist, geneticist.

Coverage Duration
Other Criteria

Plan Year

Renewal for pediatric GHD, TS, SGA, and adult GHD: patient is experiencing

improvement.

Prior Authorization Group

Drug Names

HAEGARDA HAEGARDA

Covered Uses

All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria

Required Medical Information

This medication is being used for the prevention of acute angioedema attacks. Patient has hereditary angioedema (HAE) with C1 inhibitor deficiency confirmed by laboratory testing OR patient has hereditary angioedema with normal C1 inhibitor confirmed by laboratory testing. For patients with HAE with normal C1 inhibitor, EITHER 1) Patient tested positive for the F12 gene mutation OR 2) Patient has a family history of angioedema and the angioedema was refractory to a trial of antihistamine for at least one month.

Age Restrictions

Prescriber Restrictions

Coverage Duration

Plan Year

Other Criteria

Prior Authorization Group HARVONI
Drug Names HARVONI

Covered Uses All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria -

Required Medical Information Chronic hepatitis C infection confirmed by presence of HCV RNA in the serum prior to

starting treatment. Planned treatment regimen, genotype, prior treatment history, presence or absence of cirrhosis (compensated or decompensated [Child Turcotte Pugh class B or C]), presence or absence of HIV coinfection, presence or absence of resistance-associated substitutions where applicable, liver and kidney transplantation status if applicable. Coverage conditions and specific durations of approval will be

based on current AASLD treatment guidelines.

Age Restrictions -

Prescriber Restrictions -

Coverage Duration Criteria applied consistent with current AASLD-IDSA guidance.Reminder for 8wk option

if appropriate.

Other Criteria -

Prior Authorization Group HERCEPTIN Drug Names HERCEPTIN

Covered Uses All FDA-approved indications not otherwise excluded from Part D, neoadjuvant

treatment for HER2-positive breast cancer, recurrent HER2-positive breast cancer, leptomeningeal metastases from HER2-positive breast cancer, HER2-positive

esophageal and esophagogastric junction cancer.

Exclusion Criteria -

Required Medical Information -

Age Restrictions -

Prescriber Restrictions -

Coverage Duration Neoadjuvant therapy for breast cancer: 6 months. Other: Plan Year.

Other Criteria Coverage under Part D will be denied if coverage is available under Part A or Part B as

the medication is prescribed and dispensed or administered for the individual.

Prior Authorization Group HERCEPTIN HYLECTA

Drug Names HERCEPTIN HYLECTA

Covered Uses All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria -

Required Medical Information Age Restrictions -

Prescriber Restrictions -

Coverage Duration Plan Year

Other Criteria Coverage under Part D will be denied if coverage is available under Part A or Part B as

the medication is prescribed and dispensed or administered for the individual.

Prior Authorization Group HETLIOZ HETLIOZ Drug Names

Covered Uses All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria

Required Medical Information For initial therapy and continuation of Hetlioz therapy: 1) diagnosis of Non-24 Hour

Sleep-Wake Disorder and 2) diagnosis of total blindness in both eyes (e.g., nonfunctioning retinas) and 3) unable to perceive light in both eyes. For patients currently on therapy with the requested medication, must meet at least one of the following: 1) increased total nighttime sleep or 2) decreased daytime nap duration.

Age Restrictions **Prescriber Restrictions**

Coverage Duration Initiation: 6 Months, Renewal: Plan Year

Other Criteria

Prior Authorization Group HIGH RISK MEDICATION

CYPROHEPTADINE HCL, DIGITEK, DIGOX, DIGOXIN, GUANFACINE ER. **Drug Names**

SCOPOLAMINE, TRANSDERM-SCOP

Covered Uses All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria

Required Medical Information

Age Restrictions **Prescriber Restrictions**

Coverage Duration Plan Year

Other Criteria This Prior Authorization requirement only applies to patients 70 years of age or older.

(The American Geriatrics Society identifies the use of this medication as potentially inappropriate in older adults, meaning it is best avoided, prescribed at reduced dosage,

or used with caution or carefully monitored.) Prescriber must acknowledge that

medication benefits outweigh potential risks for this patient.

Prior Authorization Group HRM-ANTICONVULSANTS

Drug Names PHENOBARBITAL, PHENOBARBITAL SODIUM

Covered Uses All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria

Required Medical Information Age Restrictions **Prescriber Restrictions**

Coverage Duration

Plan Year Other Criteria

This Prior Authorization requirement only applies to patients 70 years of age or older.

(The American Geriatrics Society identifies the use of this medication as potentially inappropriate in older adults, meaning it is best avoided, prescribed at reduced dosage,

or used with caution or carefully monitored.) Prescriber must acknowledge that

medication benefits outweigh potential risks for this patient.

Prior Authorization Group Drug Names

HRM-ANTIPARKINSON
BENZTROPINE MESYLATE, TRIHEXYPHENIDYL HCL, TRIHEXYPHENIDYL
HYDROCHLO

Covered Uses
Exclusion Criteria
Required Medical Information
Age Restrictions
Prescriber Restrictions
Coverage Duration

Other Criteria

All FDA-approved indications not otherwise excluded from Part D.

--

Plan Year

This Prior Authorization requirement only applies to patients 70 years of age or older. (The American Geriatrics Society identifies the use of this medication as potentially inappropriate in older adults, meaning it is best avoided, prescribed at reduced dosage, or used with caution or carefully monitored.) EPS (extrapyramidal symptoms): 1) One non-HRM alternative drug amantadine has not been tried. AND 2) The patient has a contraindication to one non-HRM alternative drug amantadine AND 3) Prescriber must acknowledge that medication benefits outweigh potential risks for this patient. OR 4) One non-HRM alternative drug amantadine has been tried. AND 5) The patient experienced an inadequate treatment response OR intolerance to one non-HRM alternative drug amantadine AND 6) Prescriber must acknowledge that medication benefits outweigh potential risks for this patient. Parkinson's: 1) Two non-HRM alternative drugs amantadine, carbidopa/levodopa, pramipexole, or ropinirole have been tried. AND 2) The patient experienced an inadequate treatment response OR intolerance to two non-HRM alternative drugs amantadine, carbidopa/levodopa, pramipexole, or ropinirole AND 3) Prescriber must acknowledge that medication benefits outweigh potential risks for this patient.

Prior Authorization Group Drug Names

HRM-HYDROXYZINE HYDROXYZINE HCL, HYDROXYZINE HYDROCHLORIDE, HYDROXYZINE PAMOATE

Covered Uses
Exclusion Criteria
Required Medical Information
Age Restrictions
Prescriber Restrictions
Coverage Duration

Other Criteria

All FDA-approved indications not otherwise excluded from Part D.

-Plan Year

> This Prior Authorization requirement only applies to patients 70 years of age or older. (The American Geriatrics Society identifies the use of this medication as potentially inappropriate in older adults, meaning it is best avoided, prescribed at reduced dosage, or used with caution or carefully monitored.) For pruritus: 1) A non-HRM alternative drug levocetirizine has not been tried AND 2) The patient has a contraindication to a non-HRM alternative drug levocetirizine AND 3) Prescriber must acknowledge that medication benefits outweigh potential risks for this patient OR 4) A non-HRM alternative drug levocetirizine has been tried. AND 5) The patient experienced an inadequate treatment response OR intolerance to a non-HRM alternative drug levocetirizine AND 6) Prescriber must acknowledge that medication benefits outweigh potential risks for this patient. For anxiety: 1) Two non-HRM alternative drugs buspirone, duloxetine, escitalopram, sertraline, or venlafaxine extended-release have been tried AND 2) The patient experienced an inadequate treatment response OR intolerance to two non-HRM alternative drugs buspirone, duloxetine, escitalogram, sertraline, or venlafaxine extended-release AND 3) Prescriber must acknowledge that medication benefits outweigh potential risks for this patient.

Drug Names Covered Uses

Exclusion Criteria

Required Medical Information

Age Restrictions

Prescriber Restrictions

Coverage Duration

Other Criteria

HRM-HYDROXYZINE INJ

HYDROXYZINE HCL, HYDROXYZINE HYDROCHLORIDE

All FDA-approved indications not otherwise excluded from Part D.

Plan Year

This Prior Authorization requirement only applies to patients 70 years of age or older. (The American Geriatrics Society identifies the use of this medication as potentially inappropriate in older adults, meaning it is best avoided, prescribed at reduced dosage. or used with caution or carefully monitored.) Alcohol Withdrawal Syndrome: 1) One non-HRM alternative drug clorazepate or lorazepam has not been tried AND 2) The patient has a contraindication to one non-HRM alternative drug clorazepate or lorazepam AND 3) Prescriber must acknowledge that medication benefits outweigh potential risks for this patient OR 4) One non-HRM alternative drug clorazepate or lorazepam has been tried AND 5) The patient experienced an inadequate treatment response OR intolerance to one non-HRM alternative drug clorazepate or lorazepam AND 6) Prescriber must acknowledge that medication benefits outweigh potential risks for this patient Anxiety: 1) Two non-HRM alternative drugs buspirone, duloxetine, escitalopram, sertraline or venlafaxine extended-release have been tried AND 2) The patient experienced an inadequate treatment response OR intolerance to two non-HRM alternative drugs buspirone, duloxetine, escitalopram, sertraline or venlafaxine extended-release AND 3) Prescriber must acknowledge that medication benefits outweigh potential risks for this patient OR 4) If being requested for nausea/vomiting, prescriber must acknowledge that medication benefits outweigh potential risks for this patient.

Drug Names

ZOLPIDEM TARTRATE Covered Uses All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria

Required Medical Information

Age Restrictions

Prescriber Restrictions

Coverage Duration

Other Criteria

Plan Year

HRM-HYPNOTICS

This Prior Authorization requirement only applies to patients 70 years of age or older. (The American Geriatrics Society identifies the use of this medication as potentially inappropriate in older adults, meaning it is best avoided, prescribed at reduced dosage. or used with caution or carefully monitored.) 1) One non-HRM alternative drug Silenor (3mg or 6mg) or trazodone has not been tried AND 2) The patient has a contraindication to two non-HRM alternative drugs Silenor (3mg or 6mg) and trazodone AND 3) Prescriber must acknowledge that medication benefits outweigh potential risks for this patient OR 4) One non-HRM alternative drug Silenor (3mg or 6mg) or trazodone has been tried AND 5) The patient experienced an inadequate treatment response OR intolerance to one non-HRM alternative drug Silenor (3mg or 6mg) or trazodone AND 6) Prescriber must acknowledge that medication benefits outweigh potential risks for this patient. APPLIES TO GREATER THAN CUMULATIVE 90 DAYS OF THERAPY PER YEAR.

Drug Names Covered Uses

Exclusion Criteria

Required Medical Information

Age Restrictions

Prescriber Restrictions

Coverage Duration

Other Criteria

HRM-NITROFURANTOIN

NITROFURANTOIN MACROCRYST, NITROFURANTOIN MONOHYDRAT

All FDA-approved indications not otherwise excluded from Part D.

Plan Year

This Prior Authorization requirement only applies to patients 70 years of age or older. (The American Geriatrics Society identifies the use of this medication as potentially inappropriate in older adults, meaning it is best avoided, prescribed at reduced dosage. or used with caution or carefully monitored.) 1) Two non-HRM alternative drugs cephalexin, ciprofloxacin, levofloxacin, sulfamethoxazole/trimethoprim, or trimethoprim have not been tried AND 2) The patient has a contraindication to two non-HRM alternative drugs cephalexin, ciprofloxacin, levofloxacin, sulfamethoxazole/trimethoprim, or trimethoprim AND 3) Prescriber must acknowledge that medication benefits outweigh potential risks for this patient OR 4) Two non-HRM alternative drugs cephalexin, ciprofloxacin, levofloxacin, sulfamethoxazole/trimethoprim, or trimethoprim have been tried AND 5) The patient

experienced an inadequate treatment response OR intolerance to two non-HRM alternative drugs cephalexin, ciprofloxacin, levofloxacin,

sulfamethoxazole/trimethoprim, or trimethoprim AND 6) Prescriber must acknowledge that medication benefits outweigh potential risks for this patient. APPLIES TO GREATER THAN CUMULATIVE 90 DAYS OF THERAPY PER YEAR.

Prior Authorization Group

Required Medical Information

Drug Names Covered Uses

Exclusion Criteria

Age Restrictions

Prescriber Restrictions
Coverage Duration

Other Criteria

HRM-PROMETHAZINE

PROMETHAZINE HCL, PROMETHAZINE HYDROCHLORID All FDA-approved indications not otherwise excluded from Part D.

-

-

-

-

Plan Year

This Prior Authorization requirement only applies to patients 70 years of age or older. (The American Geriatrics Society identifies the use of this medication as potentially inappropriate in older adults, meaning it is best avoided, prescribed at reduced dosage. or used with caution or carefully monitored.) Rhinitis: 1) One non-HRM alternative drug levocetirizine, azelastine nasal, fluticasone nasal, or flunisolide nasal has been tried AND 2) The patient experienced an inadequate treatment response OR intolerance to one non-HRM alternative drug levocetirizine, azelastine nasal, fluticasone nasal, or flunisolide nasal AND 3) Prescriber must acknowledge that medication benefits outweigh potential risks for this patient. Urticaria: 1) One non-HRM alternative drug levocetirizine has not been tried AND 2) The patient has a contraindication to one non-HRM alternative drug levocetirizine AND 3) Prescriber must acknowledge that medication benefits outweigh potential risks for this patient OR 4) One non-HRM alternative drug levocetirizine has been tried AND 5) The patient experienced an inadequate treatment response OR intolerance to one non-HRM alternative drug levocetirizine AND 6) Prescriber must acknowledge that medication benefits outweigh potential risks for this patient OR 7) The drug is being requested for antiemetic therapy in postoperative patients or motion sickness AND 8) Prescriber must acknowledge that medication benefits outweigh potential risks for this patient.

Prior Authorization Group

Drug Names

Covered Uses Exclusion Criteria

Required Medical Information

Age Restrictions
Prescriber Restrictions

Coverage Duration

Other Criteria

HRM-SKELETAL MUSCLE RELAXANTS CYCLOBENZAPRINE HYDROCHLO

All FDA-approved indications not otherwise excluded from Part D.

-

-Plan Year

This Prior Authorization requirement only applies to patients 70 years of age or older. (The American Geriatrics Society identifies the use of this medication as potentially inappropriate in older adults, meaning it is best avoided, prescribed at reduced dosage,

or used with caution or carefully monitored.) Prescriber must acknowledge that

medication benefits outweigh potential risks for this patient.

Prior Authorization Group Drug Names

Covered Uses

Exclusion Criteria
Required Medical Information

HUMIRA

HUMIRA, HUMIRA PEDIATRIC CROHNS D, HUMIRA PEN, HUMIRA PEN-CD/UC/HS START, HUMIRA PEN-PS/UV STARTER

All FDA-approved indications not otherwise excluded from Part D, axial spondyloarthritis.

_

For moderately to severely active rheumatoid arthritis (new starts only): 1) Inadequate response, intolerance or contraindication to methotrexate (MTX), OR 2) inadequate response or intolerance to a prior biologic disease-modifying antirheumatic drug (DMARD) or a targeted synthetic DMARD (e.g., tofacitinib). For moderately to severely active polyarticular juvenile idiopathic arthritis (new starts only): 1) Inadequate response, intolerance or contraindication to MTX, 2) inadequate response or intolerance to a prior biologic disease-modifying antirheumatic drug (DMARD). For active ankylosing spondylitis and axial spondyloarthritis (new starts only): 1) Inadequate response to a non-steroidal anti-inflammatory drug (NSAID) trial OR 2) intolerance or contraindication to NSAIDs. For moderate to severe chronic plaque psoriasis (new starts only): 1) At least 5% of body surface area (BSA) is affected OR crucial body areas (e.g., feet, hands, face, neck, groin, intertriginous areas) are affected at the time of diagnosis, AND 2) Patient meets any of the following: a) Patient has experienced an inadequate response or intolerance to either phototherapy (e.g., UVB, PUVA) or pharmacologic treatment with methotrexate, cyclosporine, or acitretin. b) Pharmacologic treatment with methotrexate, cyclosporine, or acitretin is contraindicated, or c) Patient has severe psoriasis that warrants a biologic DMARD as first-line therapy. For moderately to severely active Crohn's disease (new starts only): 1) Inadequate response to at least one conventional therapy (e.g., corticosteroids, sulfasalazine, azathioprine, mesalamine), OR 2) intolerance or contraindication to conventional therapy. For moderately to severely active ulcerative colitis (new starts only): 1) Inadequate response to at least one immunosuppressant therapy (e.g., corticosteroids, azathioprine, mercaptopurine), OR 2) intolerance or contraindication to immunosuppressant therapy.

Age Restrictions
Prescriber Restrictions
Coverage Duration
Other Criteria

-

Plan Year

_

Prior Authorization Group HYPNOTIC BENZODIAZEPINES

Drug Names TEMAZEPAM

Covered Uses All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria -

Required Medical Information Age Restrictions -

Prescriber Restrictions -

Other Criteria

Coverage Duration Plan Year

This Prior Authorization requirement only applies to patients 65 years of age or older. (The American Geriatrics Society identifies the use of this medication as potentially inappropriate in older adults, meaning it is best avoided, prescribed at reduced dosage, or used with caution or carefully monitored.) 1) One non-HRM alternative drug Silenor (3mg or 6mg) or trazodone has not been tried AND 2) The patient has a contraindication to two non-HRM alternative drugs Silenor (3mg or 6mg) and trazodone

contraindication to two non-HRM alternative drugs Silenor (3mg or 6mg) and trazodone AND 3) Prescriber must acknowledge that medication benefits outweigh potential risk in a patient 65 years of age or older OR 4) One non-HRM alternative drug Silenor (3mg or 6mg) or trazodone has been tried AND 5) The patient experienced an inadequate treatment response OR intolerance to one non-HRM alternative drug Silenor (3mg or 6mg) or trazodone AND 6) Prescriber must acknowledge that medication benefits outweigh potential risk in a patient 65 years of age or older. APPLIES TO GREATER

THAN CUMULATIVE 90 DAYS OF THERAPY PER YEAR.

Prior Authorization Group IBRANCE
Drug Names IBRANCE

Covered Uses All FDA-approved indications not otherwise excluded from Part D,

well-differentiated/dedifferentiated liposarcoma.

Exclusion Criteria - Required Medical Information -

Age Restrictions - Prescriber Restrictions -

Coverage Duration Plan Year

Other Criteria -

Prior Authorization GroupICLUSIGDrug NamesICLUSIG

Covered Uses All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria -

Required Medical Information For chronic myeloid leukemia (CML) or Philadelphia chromosome positive acute

lymphoblastic leukemia (Ph+ ALL), diagnosis was confirmed by detection of the

Philadelphia chromosome or BCR-ABL gene.

Age Restrictions Prescriber Restrictions -

Coverage Duration Plan Year

Other Criteria -

Prior Authorization Group IDHIFA
Drug Names IDHIFA

Covered UsesAll FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria Required Medical Information Age Restrictions Prescriber Restrictions -

Coverage Duration Plan Year

Other Criteria -

Prior Authorization Group IMATINIB

Drug Names IMATINIB MESYLATE

Covered Uses All FDA-approved indications not otherwise excluded from Part D, desmoid tumors,

pigmented villonodular synovitis/tenosynovial giant cell tumor (PVNS/TGCT),

chordoma, and melanoma.

Exclusion Criteria

Required Medical Information For chronic myeloid leukemia (CML) or Philadelphia chromosome positive acute

lymphoblastic leukemia (Ph+ ALL), diagnosis was confirmed by detection of the Philadelphia chromosome or BCR-ABL gene. For CML, patient did not fail (excluding failure due to intolerance) prior therapy with a tyrosine kinase inhibitor (eg, dasatinib,

nilotinib, bosutinib, ponatinib). For melanoma, c-Kit mutation is positive.

Age Restrictions -

Prescriber Restrictions -

Coverage Duration Plan Year

Other Criteria -

Prior Authorization Group IMBRUVICA
Drug Names IMBRUVICA

Covered UsesAll FDA-approved indications not otherwise excluded from Part D, gastric

mucosa-associated lymphoid tissue (MALT) lymphoma, non-gastric MALT lymphoma,

hairy cell leukemia, and lymphoplasmacytic lymphoma.

Exclusion Criteria -

Required Medical Information For mantle cell lymphoma: 1) the requested medication will be used in a patient who

has received at least one prior therapy, OR 2) the requested medication will be used in combination with rituximab as pretreatment to induction therapy with RHyperCVAD (cyclophosphamide, vincristine, doxorubicin, and dexamethasone) regimen. For gastric MALT lymphoma and non-gastric MALT lymphoma: 1) disease is recurrent, refractory, or progressive, AND 2) the requested medication will be used as second-line or subsequent therapy. For hairy cell leukemia: the requested medication will be used as

a single agent for disease progression.

Age Restrictions -

Prescriber Restrictions

Coverage Duration Plan Year

Other Criteria -

Prior Authorization Group INCRELEX
Drug Names INCRELEX

Covered Uses All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria -

Required Medical Information Must meet all of the following prior to beginning therapy with the requested medication

(new starts only): 1) height 3 or more standard deviations below the mean for children of the same age and gender AND 2) basal IGF-1 level 3 or more standard deviations below the mean for children of the same age and gender AND 3) provocative growth hormone test showing a normal or elevated growth hormone level. For renewal, patient

is experiencing improvement.

Age Restrictions -

Prescriber Restrictions -

Coverage Duration Plan Year

Other Criteria -

Prior Authorization Group INLYTA
Drug Names INLYTA

Covered UsesAll FDA-approved indications not otherwise excluded from Part D, papillary, Hurthle

cell, or follicular thyroid carcinoma.

Exclusion Criteria

Required Medical Information

For renal cell carcinoma: The disease is relapsed, metastatic, or unresectable.

Age Restrictions
Prescriber Restrictions

Coverage Duration Plan Year

Other Criteria -

Prior Authorization Group

Drug Names

IR BEFORE ER

HYSINGLA ER, METHADONE HCL, METHADONE HCL INTENSOL, MORPHINE

SULFATE ER, NUCYNTA ER

Covered Uses All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria -

Required Medical Information 1) The requested drug is being prescribed for pain associated with cancer, sickle cell

disease, a terminal condition, or pain being managed through palliative care OR 2) The

requested drug is being prescribed for pain severe enough to require daily,

around-the-clock, long-term treatment in a patient who has been taking an opioid [Note:

This drug should be prescribed only by healthcare professionals who are

knowledgeable in the use of potent opioids for the management of chronic pain.] AND 3) The patient can safely take the requested dose based on their history of opioid use

AND 4) The patient has been evaluated and will be monitored for the development of opioid use disorder AND 5) The request is for continuation of therapy for a patient who has been receiving an extended-release opioid agent for at least 30 days OR 6) The patient has severe continuous pain and has received an immediate-release opioid for

at least one week

Age Restrictions --

Coverage Duration Plan Year

Other Criteria -

Prior Authorization Group IRESSA
Drug Names IRESSA

Covered Uses All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria -

Required Medical Information For non-small cell lung cancer, patient has a known sensitizing EGFR mutation.

Age Restrictions - Prescriber Restrictions -

Coverage Duration Plan Year

Other Criteria -

Prior Authorization Group

Drug Names Covered Uses **ISOTRETINOIN**

AMNESTEEM, CLARAVIS, ISOTRETINOIN, MYORISAN, ZENATANE

All FDA-approved indications not otherwise excluded from Part D, refractory acne, severe refractory rosacea, neuroblastoma, cutaneous T-cell lymphoma (CTCL) (e.g., mycosis fungoides, Sezary syndrome), high risk for developing skin cancer (squamous cell cancers), transient acantholytic dermatosis (Grover's Disease), keratosis follicularis

(Darier Disease), lamellar ichthyosis, pityriasis rubra pilaris.

Exclusion Criteria

Required Medical Information

Age Restrictions

Prescriber Restrictions

Coverage Duration

Other Criteria

Prior Authorization Group

Drug Names Covered Uses ITRACONAZOLE

Plan Year

ITRACONAZOLE

All FDA-approved indications not otherwise excluded from Part D, Coccidioidomycosis,

Cryptococcosis, Microsporidiosis, Penicilliosis, Sporotrichosis, Pityriasis

versicolor/Tinea versicolor, Tinea corporis/Tinea cruris, Tinea manuum/Tinea pedis.

Exclusion Criteria

Required Medical Information

If for the treatment of onychomycosis due to tinea, the diagnosis has been confirmed

by a fungal diagnostic test.

Age Restrictions

Prescriber Restrictions

Coverage Duration

6 months

Other Criteria

_

Prior Authorization Group Drug Names

IVIG

BIVIGAM, CARIMUNE NANOFILTERED, FLEBOGAMMA DIF, GAMMAGARD LIQUID, GAMMAGARD S/D IGA LESS TH, GAMMAKED, GAMMAPLEX, GAMUNEX-C,

OCTAGAM, PANZYGA, PRIVIGEN

Covered Uses

All FDA-approved indications not otherwise excluded from Part D, primary immunodeficiency, chronic inflammatory demyelinating polyneuropathy, multifocal motor neuropathy, dermatomyositis, polymyositis, Guillain-Barre syndrome (GBS), myasthenia gravis, Lambert-Eaton myasthenic syndrome, Kawasaki syndrome, idiopathic thrombocytopenic purpura, pure red cell aplasia (PRCA), fetal/neonatal alloimmune thrombocytopenia, Stiff-person syndrome, and prophylaxis of bacterial infections in B-cell chronic lymphocytic leukemia (CLL), bone marrow/hematopoietic stem cell transplant (BMT/HSCT) recipients, and pediatric HIV infection.

Exclusion Criteria
Required Medical Information

_

For CLL: serum IgG less than 500 mg/dL OR a history of recurrent bacterial infections. For BMT/HSCT: IVIG is requested within the first 100 days post-transplant OR serum IgG less than 400 mg/dL. For pediatric HIV infection: 1) Serum IgG less than 400 mg/dL, OR 2) History of recurrent bacterial infections. For dermatomyositis and polymyositis: at least one standard first-line treatment (corticosteroids or immunosuppressants) has been tried but was unsuccessful or not tolerated OR patient is unable to receive standard therapy because of a contraindication or other clinical

reason. PRCA is secondary to parvovirus B19 infection. For pediatric HIV infection: age 12 years or younger.

Age Restrictions
Prescriber Restrictions
Coverage Duration
Other Criteria

Plan Year

Coverage under Part D will be denied if coverage is available under Part A or Part B as the medication is prescribed and dispensed or administered for the individual.

Prior Authorization Group
Drug Names
Covered Uses

JAKAFI JAKAFI

All FDA-approved indications not otherwise excluded from Part D, low-risk, intermediate-risk, accelerated phase, or blast phase myelofibrosis, polycythemia vera in patients with inadequate response or intolerance to interferon therapy (interferon alfa-2b, peginterferon alfa-2a, or peginterferon alfa-2b).

Exclusion Criteria

Required Medical Information

Age Restrictions
Prescriber Restrictions

Coverage Duration Plan Year

Other Criteria -

Prior Authorization Group
Drug Names
Covered Uses
Exclusion Criteria
Required Medical Information

JUXTAPID JUXTAPID

All FDA-approved indications not otherwise excluded from Part D.

_

For initiation of therapy: 1) Patient has a diagnosis of homozygous familial hypercholesterolemia (HoFH) confirmed by genetic analysis or clinical criteria (see Other Criteria), AND 2) Prior to initiation of treatment with the requested drug, patient is/was receiving a combination lipid-lowering regimen consisting of at least 2 of the following treatment options: high-intensity statin (eg, atorvastatin, rosuvastatin), fibrate (eg, fenofibrate, fenofibric acid, gemfibrozil), bile acid sequestrant (eg, cholestyramine, colesevelam, colestipol), ezetimibe, or niacin, at maximally tolerated doses or at the maximum doses approved by the FDA, AND 3) Prior to initiation of treatment with the requested drug, patient is/was experiencing an inadequate response to such combination regimen as demonstrated by treated LDL-C greater than 100 mg/dl (or greater than 70 mg/dL with clinical atherosclerotic cardiovascular disease). For renewal of therapy: 1) Patient meets all initial criteria AND 2) Has responded to therapy as demonstrated by a reduction in LDL-C.

Age Restrictions
Prescriber Restrictions
Coverage Duration
Other Criteria

Plan Year

Diagnosis of HoFH must be confirmed by one of the following: 1) Genetic diagnosis: Mutations in both alleles at LDL receptor, ApoB, PCSK9 or LDL receptor adaptor protein/ARH gene locus, or 2) Clinical diagnosis: Untreated LDL-C greater than 500 mg/dL or unknown untreated LDL-C with treated LDL-C greater than 300 mg/dL plus one of the following: a) Tendon or cutaneous xanthomas at age 10 or younger, or b) Diagnosis of FH by genetic analysis, Simon-Broome Diagnostic Criteria or Dutch Lipid Clinic Network Criteria in both parents, or c) Evidence of FH in both parents with a history including any of the following: Total cholesterol greater than or equal to 310 mg/dL, premature ASCVD [before 55 years in men and 60 years in women], tendon xanthoma, or sudden premature cardiac death. Diagnosis of FH must be confirmed by one of the following: 1) Genetic diagnosis: An LDL-receptor mutation, familial defective apo B-100, or a PCSK9 gain-of-function mutation, or 2) Simon-Broome Diagnostic Criteria for FH: Total cholesterol greater than 290 mg/dL or LDL-C greater than 190 mg/dL, plus tendon xanthoma in patient, first-degree (parent, sibling or child) or second-degree relative (grandparent, uncle or aunt), or family history of myocardial infarction in a first degree relative before the age 60 or in a second degree relative before age 50, or total cholesterol greater than 290 mg/dL in an adult first or second degree relative, or total cholesterol greater than 260 mg/dL in a child, brother, or sister aged younger than 16 years, or 3) Dutch Lipid Clinic Network Criteria for FH: Total score greater than 5 points.

Prior Authorization GroupKALYDECODrug NamesKALYDECO

Covered Uses All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria -

Required Medical Information The patient has one mutation in the cystic fibrosis transmembrane conductance

regulator (CFTR) gene that is responsive to ivacaftor potentiation based on clinical and/or in vitro assay data. If the patient's genotype is unknown, an FDA-cleared CF

mutation test should be used to detect the presence of a CFTR mutation.

Age Restrictions 6 months of age or older.

Prescriber Restrictions -

Coverage Duration Plan Year

Other Criteria The requested drug will not be used in combination with lumacaftor/ivacaftor or

tezacaftor/ivacaftor.

KETOCONAZOLE

Prior Authorization Group

Exclusion Criteria

Drug Names KETOCONAZOLE
Covered Uses All FDA-approved i

Covered Uses All FDA-approved indications not otherwise excluded from Part D, Cushing's syndrome.

Acute or chronic liver disease. Current use with dofetilide, quinidine, pimozide, cisapride, methadone, disopyramide, dronedarone, ranolazine, ergot alkaloids,

alprazolam or simvastatin.

Required Medical Information 1) Patient has one of the following diagnoses: blastomycosis, coccidioidomycosis,

histoplasmosis, chromomycosis, or paracoccidioidomycosis, OR 2) The requested drug is being prescribed for a patient with Cushing's syndrome who cannot tolerate surgery

or surgery has not been curative.

Age Restrictions -Prescriber Restrictions --

Coverage Duration 6 months

Other Criteria -

Prior Authorization Group KEYTRUDA Drug Names KEYTRUDA

Covered Uses All FDA-approved indications not otherwise excluded from Part D, malignant pleural

mesothelioma, Merkel cell carcinoma.

Exclusion Criteria -

Required Medical Information -

Age Restrictions -

Prescriber Restrictions -

Coverage Duration Plan Year

Other Criteria -

Prior Authorization Group KISQALI

Drug Names KISQALI, KISQALI FEMARA 200 DOSE, KISQALI FEMARA 400 DOSE, KISQALI

FEMARA 600 DOSE

Covered Uses All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria -

Required Medical Information -

Age Restrictions - Prescriber Restrictions -

Coverage Duration Plan Year

Other Criteria -

Prior Authorization GroupKORLYMDrug NamesKORLYM

Covered Uses All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria - Required Medical Information - Age Restrictions -

Prescriber Restrictions -

Coverage Duration Plan Year

Other Criteria -

Prior Authorization Group KUVAN Drug Names KUVAN

Covered Uses All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria -

Required Medical Information For patients who have not yet received a therapeutic trial of the requested drug, the

patient's pretreatment, including before dietary management, phenylalanine level is greater than 6 mg/dL (360 micromol/L). For patients who completed a therapeutic trial

of the requested drug, the patient must have experienced a reduction in blood

phenylalanine level of greater than or equal to 30 percent from baseline OR the patient has demonstrated an improvement in neuropsychiatric symptoms.

Age Restrictions

Prescriber Restrictions -

Coverage Duration Initial: 2 months. All others: Plan Year.

Other Criteria -

Prior Authorization Group
Drug Names
Covered Uses
Exclusion Criteria
Required Medical Information

KYNAMRO KYNAMRO

All FDA-approved indications not otherwise excluded from Part D.

_

For initiation of therapy: 1) Patient has a diagnosis of homozygous familial hypercholesterolemia (HoFH) confirmed by genetic analysis or clinical criteria (see Other Criteria), AND 2) Prior to initiation of treatment with the requested drug, patient is/was receiving a combination lipid-lowering regimen consisting of at least 2 of the following treatment options: high-intensity statin (eg, atorvastatin, rosuvastatin), fibrate (eg, fenofibrate, fenofibric acid, gemfibrozil), bile acid sequestrant (eg, cholestyramine, colesevelam, colestipol), ezetimibe, or niacin, at maximally tolerated doses or at the maximum doses approved by the FDA, AND 3) Prior to initiation of treatment with the requested drug, patient is/was experiencing an inadequate response to such combination regimen, as demonstrated by treated LDL-C greater than 100 mg/dl (or greater than 70 mg/dL with clinical atherosclerotic cardiovascular disease). For renewal of therapy, 1) Patient meets all initial criteria AND 2) Has responded to therapy as demonstrated by a reduction in LDL-C.

Age Restrictions
Prescriber Restrictions
Coverage Duration
Other Criteria

-Plan Year

Diagnosis of HoFH must be confirmed by one of the following: 1) Genetic diagnosis: Mutations in both alleles at LDL receptor, ApoB, PCSK9 or LDL receptor adaptor protein/ARH gene locus, or 2) Clinical diagnosis: Untreated LDL-C greater than 500 mg/dL or unknown untreated LDL-C with treated LDL-C greater than 300 mg/dL plus one of the following: a) Tendon or cutaneous xanthomas at age 10 or younger, or b) Diagnosis of FH by genetic analysis. Simon-Broome Diagnostic Criteria or Dutch Lipid Clinic Network Criteria in both parents, or c) Evidence of FH in both parents with a history including any of the following: Total cholesterol greater than or equal to 310 mg/dL, premature ASCVD [before 55 years in men and 60 years in women], tendon xanthoma, sudden premature cardiac death. Diagnosis of FH must be confirmed by one of the following: 1) Genetic diagnosis: An LDL-receptor mutation, familial defective apo B-100, or a PCSK9 gain-of-function mutation, or 2) Simon-Broome Diagnostic Criteria for FH: Total cholesterol greater than 290 mg/dL or LDL-C greater than 190 mg/dL, plus tendon xanthoma in patient, first-degree (parent, sibling or child) or second-degree relative (grandparent, uncle or aunt), or family history of myocardial infarction in a first degree relative before the age 60 or in a second degree relative before age 50, or total cholesterol greater than 290 mg/dL in an adult first or second degree relative, or total cholesterol greater than 260 mg/dL in a child, brother, or sister aged younger than 16 years, or 3) Dutch Lipid Clinic Network Criteria for FH: Total score greater than 5 points.

Prior Authorization Group

Drug Names

LENVIMA

LENVIMA 10 MG DAILY DOSE, LENVIMA 12MG DAILY DOSE, LENVIMA 14 MG

DAILY DOSE, LENVIMA 18 MG DAILY DOSE, LENVIMA 20 MG DAILY DOSE,

LENVIMA 24 MG DAILY DOSE, LENVIMA 4 MG DAILY DOSE, LENVIMA 8 MG DAILY

DOSE

Covered Uses All FDA-approved indications not otherwise excluded from Part D, medullary thyroid

carcinoma

Exclusion Criteria

Required Medical Information Age Restrictions

Prescriber Restrictions

Coverage Duration

Plan Year

Other Criteria

Prior Authorization Group

LETAIRIS

Drug Names

AMBRISENTAN

Covered Uses

All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria

Required Medical Information

Pulmonary arterial hypertension (WHO Group 1) was confirmed by right heart catheterization. For new starts only: 1) pretreatment mean pulmonary arterial pressure is greater than or equal to 25 mmHg, 2) pretreatment pulmonary capillary wedge pressure is less than or equal to 15 mmHg, and 3) pretreatment pulmonary vascular

resistance is greater than 3 Wood units.

Age Restrictions

Prescriber Restrictions

Coverage Duration

Plan Year

Other Criteria

Prior Authorization Group

LIDOCAINE PATCHES

Drug Names

LIDOCAINE

Covered Uses

All FDA-approved indications not otherwise excluded from Part D, pain associated with diabetic neuropathy, pain associated with cancer-related neuropathy (including

treatment-related neuropathy [e.g., neuropathy associated with radiation treatment or

chemotherapy]).

Exclusion Criteria

Required Medical Information

Age Restrictions

Prescriber Restrictions

Coverage Duration

Plan Year

Other Criteria

Updated 09/01/2019 48 Prior Authorization GroupLONSURFDrug NamesLONSURF

Covered Uses All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria -

Required Medical Information For colorectal cancer: The disease is unresectable advanced or metastatic. Patient has

progressed on treatment with EITHER a) FOLFOXIRI (fluorouracil, leucovorin,

oxaliplatin, and irinotecan) regimen OR b) irinotecan- AND oxaliplatin-based regimens.

Age Restrictions Prescriber Restrictions -

Coverage Duration Plan Year

Other Criteria -

Prior Authorization GroupLORBRENADrug NamesLORBRENA

Covered UsesAll FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria Required Medical Information Age Restrictions Prescriber Restrictions -

Coverage Duration Plan Year

Other Criteria -

Prior Authorization GroupLUMIZYMEDrug NamesLUMIZYME

Covered Uses All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria -

Required Medical Information Diagnosis of Pompe disease was confirmed by an enzyme assay demonstrating a

deficiency of acid alpha-glucosidase (GAA) enzyme activity or by genetic testing.

Age Restrictions -- Prescriber Restrictions --

Coverage Duration Plan Year

Other Criteria -

Prior Authorization Group Drug Names

LUPRON

Covered Uses

(3-MONTH), LUPRON DEPOT-PED (1-MONTH, LUPRON DEPOT-PED (3-MONTH All FDA-approved indications not otherwise excluded from Part D. in combination with growth hormone for children with growth failure and advancing puberty (leuprolide acetate only), breast cancer (3.75 mg and 11.25 mg only), malignant sex cord-stromal tumors (3.75 mg and 11.25 mg), epithelial ovarian cancer/fallopian tube cancer/primary peritoneal cancer (3.75 mg and 11.25 mg), preoperative use for uterine leiomyomata (3.75 mg and 11.25 mg).

LEUPROLIDE ACETATE, LUPRON DEPOT (1-MONTH), LUPRON DEPOT

Exclusion Criteria

Required Medical Information

For central precocious puberty (CPP), patients not currently receiving therapy must meet all of the following criteria: 1) Diagnosis of CPP confirmed by: a) a pubertal response to a gonadotropin releasing hormone (GnRH) agonist test OR a pubertal level of a third generation luteinizing hormone (LH) assay AND b) assessment of bone age versus chronological age, and 2) The onset of secondary sexual characteristics

occurred prior to 8 years of age for female patients OR prior to 9 years of age for male patients. For uterine fibroids, patient must meet one of the following: 1) Diagnosis of anemia (eg. hematocrit less than or equal to 30 percent and/or hemoglobin less than or equal to 10 g/dL). OR 2) the requested drug will be used prior to surgery for uterine

fibroids.

Age Restrictions

Prescriber Restrictions

Coverage Duration

CPP: Less than 12 years old if female and less than 13 years old if male.

Fibroids: 3 months (mo), max 6 mo total. Endometriosis: 6 mo, max 12 mo total.

Others: Plan Year

Other Criteria For prostate cancer: Use as neoadjuvant therapy prior to radical prostatectomy is not

approvable.

Prior Authorization Group

Drug Names

Covered Uses

Exclusion Criteria

Required Medical Information

LYNPARZA

LYNPARZA

All FDA-approved indications not otherwise excluded from Part D.

For HER2-negative, recurrent or metastatic breast cancer patient must meet both of the

following criteria: 1) patient has a deleterious or suspected deleterious germline BRCA mutation, and 2) patient has received prior treatment with chemotherapy or endocrine

therapy.

Age Restrictions

Prescriber Restrictions

Coverage Duration

Other Criteria

Plan Year

50 Updated 09/01/2019

Prior Authorization GroupLYRICA CRDrug NamesLYRICA CR

Covered Uses All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria Required Medical Information Age Restrictions Prescriber Restrictions -

Coverage Duration Plan Year

Other Criteria -

Prior Authorization GroupMAVYRETDrug NamesMAVYRET

Covered Uses All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria Decompensated cirrhosis/moderate or severe hepatic impairment (Child Turcotte Pugh

class B or C)

Required Medical Information Chronic hepatitis C infection confirmed by presence of HCV RNA in the serum prior to

starting treatment. Planned treatment regimen, genotype, prior treatment history, presence or absence of cirrhosis (compensated or decompensated [Child Turcotte Pugh class B or C]), presence or absence of HIV coinfection, presence or absence of resistance-associated substitutions where applicable, liver and kidney transplantation status if applicable. Coverage conditions and specific durations of approval will be

based on current AASLD treatment guidelines.

Age Restrictions -

Prescriber Restrictions -

Coverage Duration Criteria will be applied consistent with current AASLD-IDSA guidance.

Other Criteria -

Prior Authorization Group MEGESTROL

Drug Names MEGESTROL ACETATE

Covered Uses All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria Required Medical Information Age Restrictions Prescriber Restrictions -

Coverage Duration Plan Year

Other Criteria -

Prior Authorization Group MEKINIST **Drug Names** MEKINIST

Covered Uses All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria -

Required Medical Information For melanoma, tumor is positive for BRAF V600 activating mutation (e.g., BRAF V600E

or BRAF V600K mutation).

Age Restrictions -

Prescriber Restrictions -

Coverage Duration Plan Year

Other Criteria -

Prior Authorization Group MEKTOVI
Drug Names MEKTOVI

Covered Uses All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria - Required Medical Information - Age Restrictions -

Prescriber Restrictions -

Coverage Duration Plan Year

Other Criteria -

Prior Authorization Group MEMANTINE

Drug Names MEMANTINE HCL, MEMANTINE HYDROCHLORIDE, MEMANTINE

HYDROCHLORIDE E

Covered Uses All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria Required Medical Information Age Restrictions Prescriber Restrictions -

Coverage Duration Plan Year

Other Criteria This edit only applies to patients less than 30 years of age.

Prior Authorization GroupMYLOTARGDrug NamesMYLOTARG

Covered Uses All FDA-approved indications not otherwise excluded from Part D, acute promyelocytic

leukemia (APL).

Exclusion Criteria Required Medical Information Age Restrictions Prescriber Restrictions -

Coverage Duration Plan Year

Other Criteria -

Prior Authorization GroupNAGLAZYME **Drug Names**NAGLAZYME

Covered Uses All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria -

Required Medical Information Diagnosis of mucopolysaccharidosis VI disease was confirmed by an enzyme assay

demonstrating a deficiency of N-acetylgalactosamine 4-sulfatase (arylsulfatase B)

enzyme activity or by genetic testing.

Age Restrictions -Prescriber Restrictions --

Coverage Duration Plan Year

Other Criteria -

Prior Authorization Group NATPARA
Drug Names NATPARA

Covered Uses All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria Acute postsurgical hypoparathyroidism (within 6 months of surgery) and expected to

recover from the hypoparathyroidism.

Required Medical Information

Age Restrictions

Prescriber Restrictions

Coverage Duration Plan Year

Other Criteria -

Prior Authorization GroupNERLYNXDrug NamesNERLYNX

Covered Uses All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria -

Required Medical Information The requested medication is initiated within two years after completing adjuvant

trastuzumab based therapy.

Age Restrictions -

Prescriber Restrictions -

Coverage Duration Plan Year

Other Criteria -

Prior Authorization GroupNEUPOGENDrug NamesNEUPOGEN

Covered UsesAll FDA-approved indications not otherwise excluded from Part D, treatment of

chemotherapy-induced febrile neutropenia (FN), following chemotherapy for acute

lymphocytic leukemia (ALL), stem cell transplantation-related indications,

myelodysplastic syndromes (MDS), agranulocytosis, aplastic anemia, HIV-related

neutropenia, neutropenia related to renal transplantation.

Exclusion Criteria Use of the requested product within 24 hours prior to or following chemotherapy or

radiotherapy.

NEXAVAR

Required Medical Information For prophylaxis or treatment of myelosuppressive chemotherapy-induced FN patients

must meet all of the following: 1) Patient has a non-myeloid cancer, 2) Patient has received, is currently receiving, or will be receiving treatment with myelosuppressive

anti-cancer therapy

Age Restrictions -

Prescriber Restrictions -

Coverage Duration 6 months

Other Criteria -

Prior Authorization Group

Drug Names NEXAVAR

Covered Uses All FDA-approved indications not otherwise excluded from Part D, acute myeloid

leukemia, soft tissue sarcoma subtypes: angiosarcoma, desmoid tumors (aggressive fibromatosis), gastrointestinal stromal tumor (GIST), medullary thyroid carcinoma,

osteosarcoma, chordoma.

Exclusion Criteria

Required Medical Information For renal cell carcinoma: the patient has relapsed, metastatic, or unresectable disease.

For thyroid carcinoma: histology is follicular, papillary, Hurthle cell or medullary. For acute myeloid leukemia: 1) the disease is relapsed or refractory, and 2) the patient has

FLT3-ITD mutation-positive disease.

Age Restrictions -

Prescriber Restrictions -

Coverage Duration Plan Year

Other Criteria -

Prior Authorization Group NINLARO Drug Names NINLARO

Covered Uses All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria -

Required Medical Information The requested drug will be used in combination with lenalidomide and dexamethasone,

pomalidomide and dexamethasone, or dexamethasone therapy.

Age Restrictions -

Prescriber Restrictions -

Coverage Duration Plan Year

Other Criteria -

Prior Authorization Group NITYR **Drug Names** NITYR

Covered Uses All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria -

Required Medical Information Diagnosis of hereditary tyrosinemia type 1 is confirmed by one of the following: 1)

biochemical testing (e.g., detection of succinylacetone in urine) or 2) DNA testing

(mutation analysis).

Age Restrictions -

Prescriber Restrictions -

Coverage Duration Plan Year

Other Criteria -

Prior Authorization GroupNORTHERADrug NamesNORTHERA

Covered Uses All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria

Required Medical Information Prior to initial therapy, patient has a persistent, consistent decrease in systolic blood

pressure of at least 20 mmHg OR decrease in diastolic blood pressure of at least 10 mmHg within 3 minutes of standing. The requested drug will be used for patients with neurogenic orthostatic hypotension associated with one of the following diagnoses: 1) Primary autonomic failure due to Parkinson's disease, multiple system atrophy, or pure autonomic failure, OR 2) Dopamine beta hydroxylase deficiency, OR 3) Non-diabetic

autonomic neuropathy.

Age Restrictions -

Prescriber Restrictions - 3 months

Other Criteria Patients currently on Northera must experience a sustained decrease in dizziness.

Prior Authorization Group Drug NamesNUEDEXTA
NUEDEXTA

Covered Uses All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria Required Medical Information Age Restrictions Prescriber Restrictions -

Coverage Duration Plan Year

Other Criteria -

Prior Authorization GroupNUPLAZIDDrug NamesNUPLAZID

Covered Uses All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria Required Medical Information Age Restrictions Prescriber Restrictions -

Coverage Duration Plan Year

Other Criteria -

Prior Authorization Group OCTREOTIDE

Drug Names OCTREOTIDE ACETATE

Covered UsesAll FDA-approved indications not otherwise excluded from Part D, meningiomas,

thymomas and thymic carcinomas, and neuroendocrine tumors (NETs) of the

gastrointestinal (GI) tract, thymus, lung, pancreas and adrenal gland.

Exclusion Criteria -

Required Medical Information For acromegaly: 1) Patient has a high pretreatment insulin-like growth factor-1 (IGF-1)

level for age and/or gender, and 2) Patient had an inadequate or partial response to surgery or radiotherapy OR there is a clinical reason for why the patient has not had surgery or radiotherapy. For NETs of the GI tract, thymus, and lung: patient has metastatic or unresectable disease. For meningiomas: patient has unresectable

disease.

Age Restrictions -

Prescriber Restrictions -

Coverage Duration Plan Year

Other Criteria For acromegaly continuation of therapy: patient's IGF-1 level has decreased or

normalized since initiation of therapy.

Prior Authorization GroupODOMZODrug NamesODOMZO

Covered Uses All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria Required Medical Information Age Restrictions Prescriber Restrictions -

Coverage Duration Plan Year

Other Criteria -

Prior Authorization Group OFEV **Drug Names** OFEV

Covered Uses All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria -

Required Medical Information Initial Review Only: The patient does not have a known etiology for interstitial lung

disease and meets one of the following: 1) a high-resolution computed tomography

(HRCT) study of the chest or surgical lung biopsy reveals the usual interstitial pneumonia (UIP) pattern, or 2) HRCT study of the chest reveals a possible UIP pattern

and the diagnosis is supported either by surgical lung biopsy or by a multidisciplinary discussion between at least a radiologist and pulmonologist who are experienced in idiopathic pulmonary fibrosis if surgical lung biopsy has not been conducted. For continuation: The patient does not have a known etiology for interstitial lung disease.

Age Restrictions -

Prescriber Restrictions -

Coverage Duration Plan Year

Other Criteria -

Prior Authorization Group ONFI

Drug Names CLOBAZAM

Covered Uses All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria - Required Medical Information -

Age Restrictions 2 years of age or older

Prescriber Restrictions -

Coverage Duration Plan Year

Other Criteria -

Prior Authorization GroupOPSUMITDrug NamesOPSUMIT

Covered Uses All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria -

Required Medical Information Pulmonary arterial hypertension (WHO Group 1) was confirmed by right heart

catheterization. For new starts only: 1) pretreatment mean pulmonary arterial pressure is greater than or equal to 25 mmHg, 2) pretreatment pulmonary capillary wedge pressure is less than or equal to 15 mmHg, and 3) pretreatment pulmonary vascular

resistance is greater than 3 Wood units.

Age Restrictions Prescriber Restrictions -

Coverage Duration Plan Year

Other Criteria -

Prior Authorization Group ORAL-INTRANASAL FENTANYL

Drug NamesFENTANYL CITRATE, FENTANYL CITRATE ORAL TRA, FENTORACovered UsesAll FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria -

Required Medical Information 1) The requested drug is indicated for the treatment of breakthrough CANCER related

pain only. The requested drug is being prescribed for the management of breakthrough pain in a CANCER patient who is currently receiving around-the-clock opioid therapy for underlying CANCER pain AND 2) The ICD diagnosis code provided supports the CANCER RELATED diagnosis [Note: For drug coverage approval, ICD diagnosis code

provided MUST support the CANCER RELATED diagnosis.]

Coverage Duration Plan Year

Other Criteria -

Prior Authorization Group ORFADIN
Drug Names ORFADIN

Covered Uses All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria -

Required Medical Information Diagnosis of hereditary tyrosinemia type 1 is confirmed by one of the following: 1)

biochemical testing (e.g., detection of succinylacetone in urine) or 2) DNA testing

(mutation analysis).

Age Restrictions - Prescriber Restrictions -

Coverage Duration Plan Year

Other Criteria -

Prior Authorization GroupORKAMBIDrug NamesORKAMBI

Covered Uses All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria -

Required Medical Information The patient is positive for the F508del mutation on both alleles of the cystic fibrosis

transmembrane conductance regulator (CFTR) gene.

Age Restrictions 2 years of age or older

Prescriber Restrictions -

Coverage Duration Plan Year

Other Criteria The requested drug will not be used in combination with ivacaftor or

tezacaftor/ivacaftor.

Prior Authorization Group OXANDROLONE

Drug Names OXANDROLONE

Covered Uses All FDA-approved indications not otherwise excluded from Part D, Cachexia associated

with AIDS (HIV-wasting) or to enhance growth in patients with Turner's Syndrome.

Exclusion Criteria -

Required Medical Information

Age Restrictions

Prescriber Restrictions -

Coverage Duration 6 months

Other Criteria -

Prior Authorization Group PEGASYS

Drug Names PEGASYS, PEGASYS PROCLICK

Covered Uses All FDA-approved indications not otherwise excluded from Part D, myeloproliferative

neoplasm (essential thrombocythemia, polycythemia vera, primary myelofibrosis and

post-polycythemia vera or post-essential thrombocythemia myelofibrosis).

Exclusion Criteria -

Required Medical Information For chronic hepatitis C (CHC): CHC infection confirmed by presence of HCV RNA in

serum prior to starting treatment. Planned treatment regimen, genotype, prior treatment history, presence or absence of cirrhosis (compensated or decompensated [Child Turcotte Pugh class B or C]), presence or absence of HIV coinfection, presence or absence of resistance-associated substitutions where applicable, liver and kidney transplantation status if applicable. Coverage conditions and specific durations of

approval will be based on current AASLD-IDSA treatment guidelines.

Age Restrictions -

Prescriber Restrictions -

Coverage Duration HCV=Criteria will be applied consistent with current AASLD-IDSA guidance. HBV=48

wks. Other=Plan Yr

Other Criteria -

Prior Authorization Group PHENYLBUTYRATE

Drug Names SODIUM PHENYLBUTYRATE

Covered Uses All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria -

Required Medical Information Diagnosis of urea cycle disorder (UCD) was confirmed by enzymatic, biochemical or

genetic testing.

Age Restrictions -

Prescriber Restrictions -

Coverage Duration Plan Year

Other Criteria -

Prior Authorization Group PIQRAY

Drug Names PIQRAY 200MG DAILY DOSE, PIQRAY 250MG DAILY DOSE, PIQRAY 300MG

DAILY DOSE

Covered Uses All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria Required Medical Information Age Restrictions Prescriber Restrictions -

Coverage Duration Plan Year

Other Criteria -

Prior Authorization GroupPOMALYSTDrug NamesPOMALYST

Covered UsesAll FDA-approved indications not otherwise excluded from Part D, systemic light chain

amyloidosis.

Exclusion Criteria -

Required Medical Information Multiple myeloma: The patient has previously received at least two prior therapies for

multiple myeloma, including an immunomodulatory agent (i.e., thalidomide,

lenalidomide) AND a proteasome inhibitor (ie, bortezomib, carfilzomib, ixazomib).

Age Restrictions -

Prescriber Restrictions -

Coverage Duration Plan Year

Other Criteria -

Prior Authorization Group PRALUENT PRALUENT Drug Names

Covered Uses All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria Required Medical Information Age Restrictions **Prescriber Restrictions**

Coverage Duration Plan Year

Other Criteria

PROMACTA Prior Authorization Group PROMACTA Drug Names

Covered Uses All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria

Required Medical Information For chronic or persistent immune thrombocytopenia (ITP): 1) For new starts: a) patient

has had an inadequate response or is intolerant to corticosteroids, immunoglobulins or splenectomy, AND b) untransfused platelet count at any point prior to the requested medication is less than 30,000/mcL OR 30,000-50,000/mcL with symptomatic bleeding or risk factor(s) for bleeding. 2) For continuation of therapy, platelet (plt) count response to the requested drug: a) current plt count is less than or equal to 200,000/mcL OR b) current plt count is greater than 200,000/mcL and dosing will be adjusted to a plt count sufficient to avoid clinically important bleeding. For thrombocytopenia associated with chronic hepatitis C: 1) For new starts: the requested drug is used for initiation and maintenance of interferon-based therapy. 2) For continuation of therapy: patient is receiving interferon-based therapy. For severe aplastic anemia (AA): 1) For

continuation of therapy, plt count response to the requested drug: a) current plt count is 50,000-200,000/mcL, OR b) current plt count is less than 50,000/mcL and patient has not received appropriately titrated therapy for at least 16 weeks, OR c) current plt count is less than 50,000/mcL and patient is transfusion-independent, OR d) current plt count is greater than 200,000/mcL and dosing will be adjusted to achieve and maintain an

appropriate target plt count.

Age Restrictions Prescriber Restrictions Coverage Duration HCV:6mo, INITIAL: ITP/AA-6mo, REAUTH: 1) ITP/AA APR-Plan Yr. 2) AA IPR-16wks

Other Criteria APR: adequate platelet response (greater than 50k/mcL), IPR: inadequate platelet

response (less than 50k/mcL)

Updated 09/01/2019 61 Prior Authorization GroupPULMOZYMEDrug NamesPULMOZYME

Covered Uses All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria -

Required Medical Information

Diagnosis of cystic fibrosis was confirmed by appropriate diagnostic or genetic testing.

Age Restrictions - Prescriber Restrictions -

Coverage Duration Plan Year

Other Criteria Coverage under Part D will be denied if coverage is available under Part A or Part B as

the medication is prescribed and dispensed or administered for the individual.

Prior Authorization GroupQUININE SULFATEDrug NamesQUININE SULFATE

Covered UsesAll FDA-approved indications not otherwise excluded from Part D, Babesiosis,

uncomplicated Plasmodium vivax malaria.

Exclusion Criteria Required Medical Information Age Restrictions Prescriber Restrictions -

Coverage Duration 1 month
Other Criteria -

Prior Authorization GroupREGRANEXDrug NamesREGRANEX

Covered Uses All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria -

Required Medical Information For the treatment of lower extremity diabetic neuropathic ulcers that extend into the

subcutaneous tissue or beyond and have an adequate blood supply.

Coverage Duration 20 weeks

Other Criteria -

Prior Authorization GroupRELISTOR INJDrug NamesRELISTOR

Covered Uses All FDA-approved indications not otherwise excluded from Part D.

Required Medical Information

Exclusion Criteria

1) The requested drug is being prescribed for opioid-induced constipation in an adult patient with advanced illness or pain caused by active cancer who requires opioid dosage escalation for palliative care OR 2) The requested drug is being prescribed for opioid-induced constipation in an adult patient with chronic non-cancer pain, including chronic pain related to prior cancer or its treatment who does not require frequent (e.g., weekly) opioid dosage escalation AND 3) The patient is unable to tolerate oral medications OR 4) An oral drug indicated for opioid-induced constipation in an adult patient with chronic non-cancer pain has been tried. (Note: Examples are Amitiza or Movantik) AND 5) The patient experienced an inadequate treatment response or intolerance to an oral drug indicated for opioid-induced constipation in an adult patient with chronic non-cancer pain. (Note: Examples are Amitiza or Movantik) OR 6) The patient has a contraindication to an oral drug indicated for opioid-induced constipation in an adult patient with chronic non-cancer pain (Note: Examples are Amitiza or Movantik).

Age Restrictions Prescriber Restrictions -

Coverage Duration 4 Months

Other Criteria -

Prior Authorization Group
Drug Names
Covered Uses

REMICADE REMICADE

All FDA-approved indications not otherwise excluded from Part D, axial spondyloarthritis, Behcet's syndrome, granulomatosis with polyangiitis (Wegener's granulomatosis), hidradenitis suppurativa, juvenile idiopathic arthritis, pyoderma gangrenosum, sarcoidosis, Takayasu's arteritis, uveitis.

Exclusion Criteria
Required Medical Information

_

For moderately to severely active Crohn's disease (new starts only): 1) Patient has fistulizing disease OR 2) inadequate response or intolerance to a self-injectable tumor necrosis factor (TNF) inhibitor (e.g., adalimumab). For moderately to severely active ulcerative colitis (new starts only): 1) Inadequate response to at least one conventional therapy (e.g., corticosteroids, sulfasalazine, azathioprine, mesalamine) OR 2) intolerance or contraindication to conventional therapy. For moderately to severely active rheumatoid arthritis (new starts only): 1) Will be used in combination with methotrexate (MTX) or leflunomide OR patient has intolerance or contraindication to MTX or leflunomide AND 2) inadequate response or intolerance to a self-injectable tumor necrosis factor (TNF) inhibitor (e.g., adalimumab) or a targeted synthetic disease-modifying antirheumatic drug (DMARD) (e.g., tofacitinib). For active ankylosing spondylitis and axial spondyloarthritis (new starts only): 1) Inadequate response to a non-steroidal anti-inflammatory drug (NSAID) trial OR 2) intolerance or contraindication to NSAIDs. For moderate to severe chronic plaque psoriasis (new starts only): 1) At least 5% of body surface area (BSA) is affected OR crucial body areas (e.g., feet, hands, face, neck, groin, intertriginous areas) are affected at time of diagnosis, AND 2) inadequate response or intolerance to a self-injectable tumor necrosis factor (TNF) inhibitor (e.g., adalimumab). For juvenile idiopathic arthritis (new starts only): Inadequate response or intolerance to a self-injectable tumor necrosis factor (TNF) inhibitor (e.g., adalimumab). For hidradenitis suppurativa (new starts only); patient has severe, refractory disease. For uveitis (new starts only): Patient has experienced an inadequate response or intolerance or has a contraindication to a trial of immunosuppressive therapy for uveitis (e.g., methotrexate, azathioprine, or mycophenolate mofetil).

Age Restrictions
Prescriber Restrictions
Coverage Duration

-

Plan Year

Other Criteria

Prior Authorization Group
Drug Names
Covered Uses

REVLIMID REVLIMID

All FDA-approved indications not otherwise excluded from Part D, systemic light chain amyloidosis, classical Hodgkin lymphoma, myelofibrosis-associated anemia, non-Hodgkin's lymphoma with the following subtypes: chronic lymphocytic leukemia/small lymphocytic lymphoma, AIDS-related diffuse large B-cell lymphoma, primary effusion lymphoma, lymphoma associated with Castleman's disease, diffuse large B-cell lymphoma, follicular lymphoma, nongastric/gastric MALT lymphoma, primary cutaneous B-cell lymphoma, splenic/nodal marginal zone lymphoma, multicentric Castleman's disease, adult T-cell leukemia/lymphoma, mycosis fungoides/Sezary syndrome, angioimmunoblastic T-cell lymphoma, peripheral T-cell lymphoma not otherwise specified, enteropathy-associated T-cell lymphoma and primary cutaneous anaplastic large cell lymphoma.

Exclusion Criteria
Required Medical Information

Myelodysplastic syndrome (MDS): Low- to intermediate-1 risk MDS with symptomatic

anemia

Coverage Duration Plan Year

Other Criteria -

Prior Authorization Group
Drug Names
Covered Uses

RITUXAN RITUXAN

All FDA-approved indications not otherwise excluded from Part D, non-Hodgkin's lymphoma subtypes [small lymphocytic lymphoma (SLL), mantle cell lymphoma, marginal zone lymphomas (nodal, splenic, MALT), Burkitt lymphoma, primary cutaneous B-cell lymphoma, Castleman's disease, AIDS-related B-cell lymphoma, hairy cell leukemia, post-transplant lymphoproliferative disorder (PTLD), lymphoblastic lymphoma], refractory immune or idiopathic thrombocytopenic purpura (ITP), autoimmune hemolytic anemia, Waldenstrom's macroglobulinemia/lymphoplasmacytic lymphoma, chronic graft-versus-host disease (GVHD), Sjogren syndrome, thrombotic thrombocytopenic purpura, refractory myasthenia gravis, Hodgkin's lymphoma (nodular lymphocyte-predominant), primary CNS lymphoma, leptomeningeal metastases from lymphomas, acute lymphoblastic leukemia, prevention of Epstein-Barr virus (EBV)-related PTLD, multiple sclerosis.

Exclusion Criteria
Required Medical Information

For moderately to severely active rheumatoid arthritis (new starts only): 1) The requested medication is used in combination with methotrexate (MTX) unless MTX is contraindicated or not tolerated AND 2) Patient has an inadequate response, intolerance or contraindication to a self-injectable tumor necrosis factor (TNF) inhibitor (e.g., adalimumab) or a targeted synthetic disease-modifying antirheumatic drug (DMARD) (e.g., tofacitinib). Hematologic malignancies must be CD20-positive. For Wegener's Granulomatosis (WG) and Microscopic Polyangiitis (MPA): The requested medication will be used in combination with glucocorticoids. For multiple sclerosis: 1) Patient has a diagnosis of relapsing remitting multiple sclerosis and 2) Patient has had an inadequate response to two or more disease-modifying drugs indicated for multiple sclerosis despite adequate duration of treatment.

Age Restrictions
Prescriber Restrictions

Coverage Duration Plan Year

Other Criteria -

Prior Authorization Group RITUXAN HYCELA RITUXAN HYCELA

Covered Uses All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria

rituximab product by intravenous infusion without experiencing severe adverse

reactions.

Age Restrictions -

Prescriber Restrictions -

Coverage Duration Plan Year

Other Criteria -

Prior Authorization GroupRUBRACADrug NamesRUBRACA

Covered Uses All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria Required Medical Information Age Restrictions Prescriber Restrictions -

Coverage Duration Plan Year

Other Criteria -

Prior Authorization Group RYDAPT Drug Names RYDAPT

Covered Uses All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria -

Required Medical Information For newly diagnosed FLT3 mutation-positive acute myeloid leukemia (AML), the

requested medication is/was used in combination with standard cytarabine with

daunorubicin or idarubicin induction followed by cytarabine consolidation

chemotherapy.

Age Restrictions 18 years of age or older

Prescriber Restrictions -

Coverage Duration Plan Year

Other Criteria -

Prior Authorization GroupSIGNIFORDrug NamesSIGNIFOR

Covered Uses All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria -

Required Medical Information Patient has had pituitary surgery that was not curative or the patient is not a candidate

for surgery.

Age Restrictions -

Prescriber Restrictions -

Coverage Duration Plan Year

Other Criteria -

Prior Authorization Group SILDENAFIL

Drug Names SILDENAFIL CITRATE

Covered Uses All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria -

Required Medical Information Pulmonary arterial hypertension (WHO Group 1) was confirmed by right heart

catheterization. For new starts only: 1) pretreatment mean pulmonary arterial pressure is greater than or equal to 25 mmHg, 2) pretreatment pulmonary capillary wedge pressure is less than or equal to 15 mmHg, and 3) pretreatment pulmonary vascular

resistance is greater than 3 Wood units.

Age Restrictions -

Prescriber Restrictions

Coverage Duration Plan Year

Other Criteria -

Prior Authorization Group SIRTURO

Drug Names SIRTURO

Covered Uses All FDA-approved indications not otherwise excluded from Part D.

SOMATULINE DEPOT

Exclusion Criteria The requested drug is being prescribed for the treatment of latent infection due to

Mycobacterium tuberculosis, drug-sensitive tuberculosis, extra-pulmonary tuberculosis,

or infection caused by the non-tuberculous mycobacteria

Required Medical Information

Age Restrictions Prescriber Restrictions -

Coverage Duration 6 Months

Other Criteria -

Prior Authorization Group

Drug Names SOMATULINE DEPOT

Covered Uses All FDA-approved indications not otherwise excluded from Part D, neuroendocrine

tumors (NETs) of the gastrointestinal (GI) tract, thymus, lung, pancreas, and adrenal

gland.

Exclusion Criteria -

Required Medical Information For acromegaly: 1) Patient has a high pretreatment insulin-like growth factor-1 (IGF-1)

level for age and/or gender, and 2) Patient had an inadequate or partial response to surgery or radiotherapy OR there is a clinical reason for why the patient has not had surgery or radiotherapy. For NETs of the GI tract, thymus, and lung: patient has

metastatic or unresectable disease.

Age Restrictions -

Prescriber Restrictions -

Coverage Duration Plan Year

Other Criteria For acromegaly continuation of therapy: patient's IGF-1 level has decreased or

normalized since initiation of therapy.

Prior Authorization Group SOMAVERT SOMAVERT Drug Names

Covered Uses All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria

Required Medical Information Patient meets both of the following criteria: 1) Patient has a high pretreatment

> insulin-like growth factor-1 (IGF-1) level for age and/or gender, and 2) Patient had an inadequate or partial response to surgery or radiotherapy OR there is a clinical reason

for why the patient has not had surgery or radiotherapy.

Age Restrictions

Prescriber Restrictions

Coverage Duration Other Criteria For continuation of therapy: patient's IGF-1 level has decreased or normalized since

initiation of therapy.

Plan Year

Prior Authorization Group

SPRYCEL SPRYCEL Drug Names

Covered Uses All FDA-approved indications not otherwise excluded from Part D, gastrointestinal

stromal tumor (GIST).

Exclusion Criteria

Required Medical Information

For chronic myelogenous leukemia (CML) or acute lymphoblastic leukemia (ALL), diagnosis was confirmed by detection of the Philadelphia chromosome or BCR-ABL gene. For CML, 1) patient has received a hematopoietic stem cell transplant, OR 2) Patient has accelerated or blast phase CML, OR 3) For chronic phase CML, patient has one of the following a) patient is 21 years of age or younger, or b) high or

intermediate risk for disease progression, or c) low risk for disease progression and has experienced resistance, intolerance or toxicity to imatinib or an alternative tyrosine kinase inhibitor. If patient experienced resistance to imatinib or an alternative tyrosine kinase inhibitor for CML, patient is negative for T315I mutation. For GIST, patient must have PDGFRA D842V mutation and disease progression on imatinib, sunitinib, or

regorafenib.

Age Restrictions

Prescriber Restrictions

Coverage Duration Plan Year

Other Criteria

Updated 09/01/2019 69 Prior Authorization GroupSTIVARGADrug NamesSTIVARGA

Covered UsesAll FDA-approved indications not otherwise excluded from Part D, progressive GIST.

Exclusion Criteria -

Required Medical Information For colorectal cancer: The disease is unresectable advanced or metastatic. The patient

has progressed on treatment with EITHER a) FOLFOXIRI (fluorouracil, leucovorin, oxaliplatin, and irinotecan) regimen OR b) irinotecan- AND oxaliplatin-based regimens.

Age Restrictions --

Coverage Duration Plan Year

Other Criteria -

Prior Authorization Group SUTENT Drug Names SUTENT

Covered UsesAll FDA-approved indications not otherwise excluded from Part D, thyroid carcinoma

(follicular, papillary, Hurthle cell, or medullary), angiosarcoma, solitary fibrous tumor,

hemangiopericytoma, chordoma (bone cancer), thymic carcinoma.

Exclusion Criteria

Required Medical Information For renal cell carcinoma: Either 1) The disease is relapsed, metastatic, or unresectable,

OR 2) The patient is at high risk of disease recurrence following nephrectomy.

Age Restrictions -

Prescriber Restrictions -

Coverage Duration Plan Year

Other Criteria -

Prior Authorization Group SYLATRON Drug Names SYLATRON

Covered UsesAll FDA-approved indications not otherwise excluded from Part D, myelofibrosis,

polycythemia vera, essential thrombocythemia.

Exclusion Criteria -

Required Medical Information -

Age Restrictions -

Prescriber Restrictions -

Coverage Duration Plan Year

Other Criteria -

Prior Authorization GroupSYMDEKODrug NamesSYMDEKO

Covered Uses All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria -

Required Medical Information The patient is positive for the F508del mutation on both alleles of the cystic fibrosis

transmembrane conductase regulator (CFTR) gene or the patient has a mutation in the cystic fibrosis transmembrane conductance regulator (CFTR) gene that is responsive to tezacaftor/ivacaftor potentiation based on clinical and/or in vitro assay data. If the patient's genotype is unknown, an FDA-cleared cystic fibrosis mutation test should be

used to detect the presence of a CFTR mutation

Age Restrictions 6 years of age or older

Prescriber Restrictions

Coverage Duration Plan Year

Other Criteria Symdeko will not be used in combination with Orkambi or Kalydeco.

Prior Authorization Group SYMPAZAN Drug Names SYMPAZAN

Covered Uses All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria -

Required Medical Information -

Age Restrictions 2 years of age or older

Prescriber Restrictions -

Coverage Duration Plan Year

Other Criteria -

Prior Authorization GroupSYNRIBODrug NamesSYNRIBO

Covered Uses All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria -

Required Medical Information For chronic myeloid leukemia (CML), the patient has experienced resistance, toxicity or

intolerance to prior therapy with at least two tyrosine kinase inhibitors (TKIs) (eg,

imatinib, dasatinib, nilotinib, bosutinib, ponatinib).

Coverage Duration Plan Year

Other Criteria -

Prior Authorization Group TAFINLAR
Drug Names TAFINLAR

Covered UsesAll FDA-approved indications not otherwise excluded from Part D, brain metastases

from melanoma.

Exclusion Criteria

Required Medical Information For melanoma (including brain metastases), tumor is positive for a BRAF V600

activating mutation (e.g., BRAF V600E or BRAF V600K mutation). For NSCLC, tumor

is positive for a BRAF V600 activating mutation.

Age Restrictions -

Prescriber Restrictions

Coverage Duration Plan Year

Other Criteria -

Prior Authorization Group TAGRISSO
Drug Names TAGRISSO

Covered Uses All FDA-approved indications not otherwise excluded from Part D, EGFR

mutation-positive recurrent or metastatic non-small cell lung cancer, brain metastases if

active against primary tumor (EGFR T790M mutation-positive non-small cell lung

cancer).

Exclusion Criteria -

Required Medical Information -

Age Restrictions -

Prescriber Restrictions

Coverage Duration Plan Year

Other Criteria -

Prior Authorization Group TALZENNA
Drug Names TALZENNA

Covered Uses All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria - Required Medical Information -

Age Restrictions -

Prescriber Restrictions -

Coverage Duration Plan Year

Other Criteria -

Prior Authorization Group TARCEVA

Drug Names ERLOTINIB HYDROCHLORIDE, TARCEVA

Covered Uses All FDA-approved indications not otherwise excluded from Part D, chordoma, renal cell

carcinoma (RCC).

Exclusion Criteria -

Required Medical Information For non-small cell lung cancer, patient has a known sensitizing EGFR mutation. For

pancreatic cancer, the disease is locally advanced, unresectable, or metastatic.

Age Restrictions -

Prescriber Restrictions -

Coverage Duration Plan Year

Other Criteria -

Prior Authorization Group TASIGNA

Drug Names TASIGNA

Covered Uses All FDA-approved indications not otherwise excluded from Part D, Philadelphia

chromosome positive acute lymphoblastic leukemia (Ph+ ALL), gastrointestinal stromal

tumor (GIST).

Exclusion Criteria -

Required Medical Information For CML or ALL, diagnosis was confirmed by detection of the Philadelphia

chromosome or BCR-ABL gene. For CML, 1) patient has received a hematopoietic stem cell transplant, OR 2) patient has accelerated or blast phase CML, OR 3) For chronic phase CML, the patient has one of the following: a) patient is 18 years of age or younger, b) high or intermediate risk for disease progression, or c) low risk for disease progression and has experienced resistance, intolerance or toxicity to imatinib or an alternative tyrosine kinase inhibitor. If patient experienced resistance to imatinib or an alternative tyrosine kinase inhibitor for CML, patient is negative for T315I mutation. For

GIST, patient must have progressed on imatinib, sunitinib or regorafenib.

Age Restrictions -

Prescriber Restrictions -

Coverage Duration Plan Year

Other Criteria -

Prior Authorization Group TAZAROTENE

Drug Names TAZAROTENE, TAZORAC

Covered Uses All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria -

Required Medical Information For plaque psoriasis, the requested drug is being prescribed to treat less than 20

percent of the patient's body surface area.

Age Restrictions -

Prescriber Restrictions -

Coverage Duration Plan Year

Other Criteria -

Prior Authorization GroupTECENTRIQDrug NamesTECENTRIQ

Covered Uses All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria Required Medical Information Age Restrictions Prescriber Restrictions -

Coverage Duration Plan Year

Other Criteria -

Prior Authorization GroupTESTOSTERONE CYPIONATE INJDrug NamesTESTOSTERONE CYPIONATE

Covered Uses All FDA-approved indications not otherwise excluded from Part D, Gender Dysphoria in

transgender male patients.

Exclusion Criteria -

Required Medical Information 1) Request is for continuation of testosterone therapy and requested drug is being

prescribed for hypogonadism in a male patient or a patient that self-identifies as male

who has had a confirmed low testosterone level according to current practice guidelines or your standard male lab reference values before starting testosterone therapy OR 2) Request is not for continuation of testosterone therapy and requested

drug is being prescribed for hypogonadism in a male patient or a patient that

self-identifies as male who has at least two confirmed low testosterone levels according

to current practice guidelines or your standard male lab reference values OR 3)

Requested drug is being prescribed for gender dysphoria in a transgender male patient who is 12 years of age or older and able to make an informed, mature decision to

engage in therapy.

Age Restrictions 12 years of age or older (applies to gender dysphoria only)

Prescriber Restrictions

Coverage Duration Plan Year

Other Criteria -

Prior Authorization Group

Drug Names

Covered Uses

Exclusion Criteria
Required Medical Information

TESTOSTERONE ENANTHATE INJ
TESTOSTERONE ENANTHATE

All FDA-approved indications not otherwise excluded from Part D.

_

1) Request is for continuation of testosterone therapy and requested drug is being prescribed for hypogonadism in a male patient or a patient that self-identifies as male who had a confirmed low testosterone level according to current practice guidelines or your standard male lab reference values before starting testosterone therapy OR 2) Request is not for continuation of testosterone therapy and requested drug is being prescribed for hypogonadism in a male patient or a patient that self-identifies as male who has at least two confirmed low testosterone levels according to current practice guidelines or your standard male lab reference values OR 3) Requested drug is being prescribed for inoperable metastatic breast cancer in a patient who is 1 to 5 years postmenopausal and who has had an incomplete response to other therapy for metastatic breast cancer OR 4) Requested drug is being prescribed for a pre-menopausal patient with breast cancer who has benefited from oophorectomy and is considered to have a hormone-responsive tumor OR 5) Requested drug is being prescribed for delayed puberty in a male patient.

Age Restrictions Prescriber Restrictions -

Coverage Duration Plan Year

Other Criteria -

Prior Authorization Group TETRABENAZINE
Drug Names TETRABENAZINE

Covered Uses All FDA-approved indications not otherwise excluded from Part D, chronic tics, tardive

dyskinesia, hemiballismus, chorea not associated with Huntington's disease.

Exclusion Criteria -

Required Medical Information -

Age Restrictions -

Prescriber Restrictions -

Coverage Duration Plan Year

Other Criteria -

Prior Authorization GroupTHALOMIDDrug NamesTHALOMID

Covered Uses All FDA-approved indications not otherwise excluded from Part D, myelofibrosis-related

anemia, systemic light chain amyloidosis, Waldenstrom's

macroglobulinemia/lymphoplasmacytic lymphoma, recurrent aphthous stomatitis, recurrent HIV-associated aphthous ulcers, cachexia, HIV-associated diarrhea, Kaposi's sarcoma, Behcet's syndrome, chronic graft-versus-host disease, Crohn's disease,

multicentric Castleman's disease.

Exclusion Criteria

Required Medical Information Cachexia: Cachexia must be due to cancer or HIV infection. Kaposi's sarcoma: The

patient has HIV infection.

Age Restrictions -

Prescriber Restrictions -

Coverage Duration Plan Year

Other Criteria -

Prior Authorization GroupTIBSOVODrug NamesTIBSOVO

Covered Uses All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria Required Medical Information Age Restrictions Prescriber Restrictions -

Coverage Duration Plan Year

Other Criteria -

Prior Authorization GroupTOBRAMYCINDrug NamesTOBRAMYCIN

Covered Uses All FDA-approved indications not otherwise excluded from Part D, non-cystic fibrosis

bronchiectasis.

Exclusion Criteria -

Required Medical Information Pseudomonas aeruginosa is present in the patient's airway cultures OR the patient has

a history of pseudomonas aeruginosa infection or colonization in the airways.

Age Restrictions -

Prescriber Restrictions -

Coverage Duration Plan Year

Other Criteria Coverage under Part D will be denied if coverage is available under Part A or Part B as

the medication is prescribed and dispensed or administered for the individual.

Prior Authorization Group

Drug Names

TOPICAL LIDOCAINE

GLYDO, LIDOCAINE, LIDOCAINE HCL, LIDOCAINE HCL JELLY.

LIDOCAINE/PRILOCAINE

Covered Uses

All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria

Required Medical Information

1) The requested drug is being used for topical anesthesia, 2) If the requested drug will

be used as part of a compounded product, then all the active ingredients in the

compounded product are FDA-approved for topical use

Age Restrictions

Prescriber Restrictions

Coverage Duration

Other Criteria

3 Months

Coverage under Part D will be denied if coverage is available under Part A or Part B as

the medication is prescribed and dispensed or administered for the individual.

Prior Authorization Group

Drug Names Covered Uses Exclusion Criteria

Required Medical Information

TOPICAL TESTOSTERONES

ANDRODERM, TESTOSTERONE, TESTOSTERONE PUMP All FDA-approved indications not otherwise excluded from Part D.

1) Request is for continuation of testosterone therapy and requested drug is being prescribed for hypogonadism in a male patient or a patient that self-identifies as male who had a confirmed low testosterone level according to current practice guidelines or your standard male lab reference values before starting testosterone therapy OR 2) Request is not for continuation of testosterone therapy and requested drug is being prescribed for hypogonadism in a male patient or a patient that self-identifies as male who has at least two confirmed low testosterone levels according to current practice guidelines or your standard male lab reference values.

Age Restrictions

Prescriber Restrictions

Plan Year **Coverage Duration**

Other Criteria

Prior Authorization Group TOPICAL TRETINOIN **Drug Names** AVITA, TRETINOIN

Covered Uses All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria Required Medical Information Age Restrictions Prescriber Restrictions

Coverage Duration Plan Year

Other Criteria

77 Updated 09/01/2019

Prior Authorization Group TRELSTAR

Drug Names TRELSTAR MIXJECT

Covered Uses All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria -

Required Medical Information -

Age Restrictions -Prescriber Restrictions --

Coverage Duration Plan Year

Other Criteria Use as neoadjuvant therapy prior to radical prostatectomy is not approvable.

Prior Authorization Group TREPROSTINIL INJ

Drug Names REMODULIN, TREPROSTINIL

Covered Uses All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria -

Required Medical Information Pulmonary arterial hypertension (WHO Group 1) was confirmed by right heart

catheterization. For new starts only: 1) pretreatment mean pulmonary arterial pressure is greater than or equal to 25 mmHg, 2) pretreatment pulmonary capillary wedge pressure is less than or equal to 15 mmHg, and 3) pretreatment pulmonary vascular

resistance is greater than 3 Wood units.

Age Restrictions -

Prescriber Restrictions -

Coverage Duration Plan Year

Other Criteria Coverage under Part D will be denied if coverage is available under Part A or Part B as

the medication is prescribed and dispensed or administered for the individual.

Prior Authorization Group TRIENTINE

Drug Names TRIENTINE HYDROCHLORIDE

Covered Uses All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria - Required Medical Information -

Age Restrictions -- Prescriber Restrictions --

Coverage Duration Plan Year

Other Criteria -

Prior Authorization Group TYKERB
Drug Names TYKERB

Covered Uses All FDA-approved indications not otherwise excluded from Part D, metastatic CNS

lesions from HER2-positive breast cancer.

Exclusion Criteria

Required Medical Information For HER2-positive breast cancer, the requested drug will be used in combination with:

1) aromatase inhibitor (e.g., anastrozole, letrozole, exemestane), or 2) capecitabine, or

3) trastuzumab.

Age Restrictions -

Prescriber Restrictions

Coverage Duration Plan Year

Other Criteria -

Prior Authorization Group TYMLOS

Drug Names TYMLOS

Covered UsesAll FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria -

Required Medical InformationFor postmenopausal osteoporosis: patient has ONE of the following (1. or 2.): 1) A history of fragility fractures, OR 2) A pre-treatment T-score of less than or equal to -2.5 or osteopenia with a high pre-treatment FRAX fracture probability and patient has ANY

of the following: a) Indicators for higher fracture risk (e.g., advanced age, frailty, glucocorticoid therapy, very low T-scores, or increased fall risk), OR b) Patient has failed prior treatment with or is intolerant to a previous osteoporosis therapy (i.e., oral

bisphosphonates or injectable antiresorptive agents)

Age Restrictions --

Coverage Duration 24 months lifetime total for parathyroid hormone analogs (e.g., abaloparatide or

teriparatide)

Other Criteria Patient has high FRAX fracture probability if the 10 year probability is either greater

than or equal to 20% for any major osteoporotic fracture or greater than or equal to 3%

for hip fracture

Prior Authorization Group VALCHLOR Drug Names VALCHLOR

Covered Uses All FDA-approved indications not otherwise excluded from Part D, chronic or

smoldering adult T-cell leukemia/lymphoma, mycosis fungoides, primary cutaneous marginal zone lymphoma, primary cutaneous follicle center lymphoma, lymphomatoid

papulosis.

Exclusion Criteria -

Required Medical Information

Age Restrictions -

Prescriber Restrictions -

Coverage Duration Plan Year

Other Criteria -

Prior Authorization Group VELCADE

Drug Names BORTEZOMIB, VELCADE

Covered Uses All FDA-approved indications not otherwise excluded from Part D, systemic light chain

amyloidosis, Waldenstrom's macroglobulinemia/lymphoplasmacytic lymphoma,

multicentric Castleman's disease.

Exclusion Criteria -

Required Medical Information -

Age Restrictions -

Prescriber Restrictions -

Coverage Duration Plan Year

Other Criteria Coverage under Part D will be denied if coverage is available under Part A or Part B as

the medication is prescribed and dispensed or administered for the individual.

Prior Authorization Group VENCLEXTA

Drug Names VENCLEXTA, VENCLEXTA STARTING PACK

Covered Uses All FDA-approved indications not otherwise excluded from Part D, small lymphocytic

lymphoma, mantle cell lymphoma.

Exclusion Criteria -

Required Medical Information -

Age Restrictions -

Prescriber Restrictions -

Coverage Duration Plan Year

Other Criteria -

Prior Authorization Group VENTAVIS **Drug Names** VENTAVIS

Covered Uses All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria -

Required Medical Information Pulmonary arterial hypertension (WHO Group 1) was confirmed by right heart

catheterization. For new starts only: 1) pretreatment mean pulmonary arterial pressure is greater than or equal to 25 mmHg, 2) pretreatment pulmonary capillary wedge pressure is less than or equal to 15 mmHg, and 3) pretreatment pulmonary vascular

resistance is greater than 3 Wood units.

Age Restrictions - Prescriber Restrictions -

Coverage Duration Plan Year

Other Criteria Coverage under Part D will be denied if coverage is available under Part A or Part B as

the medication is prescribed and dispensed or administered for the individual.

Prior Authorization Group VERSACLOZ Drug Names VERSACLOZ

Covered Uses All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria Required Medical Information Age Restrictions Prescriber Restrictions -

Coverage Duration Plan Year

Other Criteria -

Prior Authorization Group VERZENIO Drug Names VERZENIO

Covered Uses All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria Required Medical Information Age Restrictions Prescriber Restrictions -

Coverage Duration Plan Year

Other Criteria -

Prior Authorization Group VIGABATRIN

Drug Names VIGABATRIN, VIGADRONE

Covered Uses All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria -

Required Medical Information For complex partial seizures (CPS): patient had an inadequate response to at least 2

alternative therapies for CPS (e.g., carbamazepine, phenytoin, levetiracetam,

topiramate, oxcarbazepine or lamotrigine).

Age Restrictions --

Coverage Duration Plan Year

Other Criteria -

Prior Authorization Group VITRAKVI
Drug Names VITRAKVI

Covered Uses All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria Required Medical Information Age Restrictions Prescriber Restrictions -

Coverage Duration Plan Year

Other Criteria -

Prior Authorization GroupVIZIMPRODrug NamesVIZIMPRO

Covered Uses All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria Required Medical Information Age Restrictions Prescriber Restrictions -

Coverage Duration Plan Year

Other Criteria -

Prior Authorization Group VOSEVI
Drug Names VOSEVI

Covered Uses All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria Decompensated cirrhosis/moderate or severe hepatic impairment (Child Turcotte Pugh

class B or C)

Required Medical Information Chronic hepatitis C infection confirmed by presence of HCV RNA in the serum prior to

starting treatment. Planned treatment regimen, genotype, prior treatment history, presence or absence of cirrhosis (compensated or decompensated [Child Turcotte Pugh class B or C]), presence or absence of HIV coinfection, presence or absence of resistance-associated substitutions where applicable, liver and kidney transplantation status if applicable. Coverage conditions and specific durations of approval will be

based on current AASLD treatment guidelines.

Age Restrictions -

Prescriber Restrictions -

Coverage Duration Criteria will be applied consistent with current AASLD-IDSA guidance.

Other Criteria -

Prior Authorization Group VOTRIENT Drug Names VOTRIENT

Covered UsesAll FDA-approved indications not otherwise excluded from Part D, thyroid carcinoma

(follicular, papillary, Hurthle cell, or medullary), uterine sarcoma, ovarian cancer

(epithelial ovarian, fallopian tube, or primary peritoneal).

Exclusion Criteria -

Required Medical Information For renal cell carcinoma: The disease is relapsed, metastatic, or unresectable. For soft

tissue sarcoma (STS): 1) The patient does not have an adipocytic soft tissue sarcoma, AND 2) The patient has one of the following subtypes of STS: a) gastrointestinal stromal tumor (GIST), b) angiosarcoma, c) pleomorphic rhabdomyosarcoma, d) retroperitoneal/intra-abdominal sarcoma, or e) extremity/superficial trunk, head/neck

sarcoma.

Age Restrictions -

Prescriber Restrictions -

Coverage Duration Plan Year

Other Criteria -

Prior Authorization GroupVRAYLARDrug NamesVRAYLAR

Covered Uses All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria -

Required Medical Information The patient experienced an inadequate treatment response, intolerance, or

contraindication to one of the following: lurasidone, aripiprazole, olanzapine,

paliperidone, quetiapine, risperidone, or ziprasidone.

Age Restrictions - Prescriber Restrictions -

Coverage Duration Plan Year

Other Criteria -

Prior Authorization Group XALKORI Drug Names XALKORI

Covered Uses All FDA-approved indications not otherwise excluded from Part D, non-small cell lung

cancer (NSCLC) with high-level MET amplification or MET exon 14 skipping mutation,

inflammatory myofibroblastic tumors (IMT).

Exclusion Criteria

Required Medical Information For IMT, the tumor is ALK-positive.

Age Restrictions -Prescriber Restrictions --

Coverage Duration Plan Year

Other Criteria -

Prior Authorization Group XELJANZ

Drug Names XELJANZ, XELJANZ XR

Covered Uses All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria -

Required Medical Information For moderately to severely active rheumatoid arthritis (new starts only): Patient meets

at least one of the following criteria: 1) Inadequate response, intolerance or

contraindication to methotrexate (MTX), OR 2) Inadequate response or intolerance to a prior biologic disease-modifying antirheumatic drug (DMARD) (e.g., adalimumab). For active psoriatic arthritis (new starts only): Patient meets BOTH of the following criteria: 1) Inadequate response to methotrexate (MTX) or other nonbiologic disease-modifying antirheumatic drugs (DMARDs) (e.g., leflunomide, sulfasalazine, etc.) OR a prior biologic DMARD (e.g., adalimumab), AND 2) The requested drug is used in combination with a nonbiologic DMARD (e.g., methotrexate, leflunomide, sulfasalazine,

etc.). For moderately to severely active ulcerative colitis (new starts only): Patient meets at least one of the following criteria: 1) Inadequate response, intolerance or contraindication to at least one conventional therapy option (e.g., oral aminosalicylates,

corticosteroids), or 2) Inadequate response or intolerance to a prior biologic disease-modifying antirheumatic drug (DMARD) (e.g., adalimumab)

Age Restrictions - Prescriber Restrictions -

Coverage Duration Plan Year

Other Criteria -

Prior Authorization Group XGEVA
Drug Names XGEVA

Covered Uses All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria -

Required Medical Information For hypercalcemia of malignancy, condition is refractory to intravenous (IV)

bisphosphonate therapy (eg, zoledronic acid, pamidronate) or there is a clinical reason

to avoid IV bisphosphonate therapy.

Age Restrictions Prescriber Restrictions -

Coverage Duration Plan Year

Other Criteria Coverage under Part D will be denied if coverage is available under Part A or Part B as

the medication is prescribed and dispensed or administered for the individual.

Prior Authorization Group XIFAXAN
Drug Names XIFAXAN

Covered Uses All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria Required Medical Information Age Restrictions Prescriber Restrictions -

Coverage Duration Reduction in risk of overt hepatic encephalopathy recurrence-6 Months. IBS-D - Plan

Year

Other Criteria -

Prior Authorization GroupXOLAIRDrug NamesXOLAIR

Covered Uses All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria

Required Medical Information For allergic asthma initial therapy: 1)Patient has positive skin test (or blood test) to at

least 1 perennial aeroallergen, 2) Patient has baseline IgE level greater than or equal to 30 IU/mL, 3) Patient has inadequate asthma control despite current treatment with both of the following medications at optimized doses: a) Inhaled corticosteroid, b) Additional controller (long acting beta2-agonist, leukotriene modifier, or sustained-release

controller (long acting beta2-agonist, leukotriene modifier, or sustained-release theophylline) unless patient has an intolerance or contraindication to such therapies. For allergic asthma continuation therapy only: Patient's asthma control has improved on the requested drug since initiation of therapy. Chronic idiopathic urticaria (CIU) initial

therapy: 1) Patient has been evaluated for other causes of urticaria, including bradykinin-related angioedema and IL-1-associated urticarial syndromes

(auto-inflammatory disorders, urticarial vasculitis), 2) Patient has experienced a spontaneous onset of wheals, angioedema, or both, for at least 6 weeks. For CIU continuation therapy: Patient has experienced a response (e.g., improved symptoms)

since initiation of therapy.

Age Restrictions For CIU: 12 years of age or older. For allergic asthma: 6 years of age or older.

Prescriber Restrictions -

Coverage Duration Allergic asthma: Plan Year. CIU initial: 6 months. CIU continuation: Plan Year.

Other Criteria -

Prior Authorization Group XOSPATA
Drug Names XOSPATA

Covered Uses All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria -

Required Medical Information -

Age Restrictions 18 years of age or older

Prescriber Restrictions -

Coverage Duration Plan Year

Other Criteria -

Prior Authorization GroupXTANDIDrug NamesXTANDI

Covered Uses All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria -

Required Medical Information The requested drug will be used to treat prostate cancer.

Age Restrictions - Prescriber Restrictions -

Coverage Duration Plan Year

Other Criteria -

Prior Authorization Group XYREM Drug Names XYREM

Covered Uses All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria -

Required Medical Information 1) The drug is being prescribed for the treatment of excessive daytime sleepiness in a

patient with narcolepsy AND 2) The patient experienced an inadequate treatment response or intolerance to at least one CNS stimulant drug and one CNS promoting wakefulness drug OR 3) the patient has a contraindication to at least one CNS stimulant drug and one CNS wakefulness promoting drug (NOTE: Examples of a CNS stimulant drug are amphetamine, dextroamphetamine, or methylphenidate. Example of

a CNS wakefulness promoting drug is armodafinil. Coverage of armodafinil or

amphetamines or methylphenidates may require prior authorization). OR 4) The drug is

being prescribed for the treatment of cataplexy in a patient with narcolepsy

Age Restrictions - Prescriber Restrictions -

Coverage Duration Plan Year

Other Criteria If the request is for the continuation of Xyrem (sodium oxybate), then the patient

experienced a decrease in daytime sleepiness with narcolepsy or a decrease in

cataplexy episodes with narcolepsy.

Prior Authorization Group ZAVESCA
Drug Names MIGLUSTAT

Covered Uses All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria -

Required Medical Information Diagnosis of Gaucher disease was confirmed by an enzyme assay demonstrating a

deficiency of beta-glucocerebrosidase enzyme activity or by genetic testing.

Age Restrictions 18 years of age or older

Prescriber Restrictions -

Coverage Duration Plan Year

Other Criteria -

Prior Authorization GroupZEJULADrug NamesZEJULA

Covered Uses All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria Required Medical Information Age Restrictions Prescriber Restrictions -

Coverage Duration Plan Year

Other Criteria Treatment is being started or was started no later than 8 weeks after the most recent

platinum-based chemotherapy.

Prior Authorization GroupZELBORAFDrug NamesZELBORAF

Covered Uses All FDA-approved indications not otherwise excluded from Part D, brain metastases

from melanoma, non-small cell lung cancer, hairy cell leukemia, and thyroid carcinoma

(papillary, follicular, and Hurthle).

Exclusion Criteria -

Required Medical Information For melanoma (including brain metastases), tumor is positive for BRAF V600 activating

mutation (e.g., BRAF V600E or BRAF V600K mutation). For non-small cell lung cancer,

tumor is positive for the BRAF V600E mutation. For thyroid carcinoma the tumor is

positive for BRAF mutation.

Age Restrictions -

Prescriber Restrictions -

Coverage Duration Plan Year

Other Criteria -

Prior Authorization Group ZEPATIER
Drug Names ZEPATIER

Covered Uses All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria Decompensated cirrhosis/moderate or severe hepatic impairment (Child Turcotte Pugh

class B or C)

Required Medical Information Chronic hepatitis C infection confirmed by presence of HCV RNA in the serum prior to

starting treatment. Planned treatment regimen, genotype, prior treatment history, presence or absence of cirrhosis (compensated or decompensated [Child Turcotte Pugh class B or C]), presence or absence of HIV coinfection, presence or absence of resistance-associated substitutions (eg, NS5A polymorphisms) where applicable, liver and kidney transplantation status if applicable. Coverage conditions and specific durations of approval will be based on current AASLD-IDSA treatment guidelines.

Age Restrictions

Prescriber Restrictions - Coverage Duration - Criteria will be applied consisten

Prior Authorization GroupZOLINZADrug NamesZOLINZA

Covered UsesAll FDA-approved indications not otherwise excluded from Part D, mycosis fungoides,

Sezary syndrome.

Exclusion Criteria -

Required Medical Information

Age Restrictions

Prescriber Restrictions -

Coverage Duration Plan Year

Other Criteria -

Prior Authorization Group ZYDELIG
Drug Names ZYDELIG

Covered Uses All FDA-approved indications not otherwise excluded from Part D, relapsed or

refractory chronic lymphocytic leukemia (CLL)/small lymphocytic lymphoma (SLL), refractory, relapsed or progressive follicular lymphoma, and marginal zone lymphomas [nodal marginal zone lymphoma, gastric mucosa associated lymphoid tissue (MALT) lymphoma, non-gastric MALT lymphoma, and splenic marginal zone lymphoma].

Exclusion Criteria -

Required Medical Information

Age Restrictions -

Prescriber Restrictions -

Coverage Duration Plan Year

Other Criteria -

Prior Authorization Group ZYKADIA
Drug Names ZYKADIA

Covered UsesAll FDA-approved indications not otherwise excluded from Part D, anaplastic lymphoma

kinase (ALK)-positive inflammatory myofibroblastic tumor.

Exclusion Criteria -

Required Medical Information For non-small cell lung cancer (NSCLC), the requested medication is used for the

treatment of recurrent or metastatic ALK-positive NSCLC. For inflammatory

myofibroblastic tumor, the tumor is ALK-positive.

Age Restrictions -

Prescriber Restrictions -

Coverage Duration Plan Year

Other Criteria -

Prior Authorization GroupZYPREXA RELPREVVDrug NamesZYPREXA RELPREVV

Covered Uses All FDA-approved indications not otherwise excluded from Part D.

Exclusion Criteria -

Required Medical Information Tolerability with oral olanzapine has been established.

Age Restrictions - Prescriber Restrictions -

Coverage Duration Plan Year

Other Criteria -

Prior Authorization Group ZYTIGA

Drug Names ABIRATERONE ACETATE, ZYTIGA

Covered Uses All FDA-approved indications not otherwise excluded from Part D and newly diagnosed

metastatic or high-risk locally advanced prostate cancer.

Exclusion Criteria -

Required Medical Information For metastatic castration-resistant prostate cancer: The requested drug will be used in

combination with prednisone. For castration-sensitive metastatic or locally advanced prostate cancer: 1) The requested drug will be used in combination with prednisone and concurrent androgen-deprivation therapy. Androgen deprivation therapy is not required in patients who have had bilateral orchiectomy, 2) Disease is newly diagnosed and metastatic, node-positive, high-risk locally advanced, or was previously treated with radical surgery or radiotherapy and is now relapsing with high risk features.

Age Restrictions -

Prescriber Restrictions -

Coverage Duration Plan Year

Other Criteria -

This prior authorization criteria is current as of 09/01/2019. For more recent information or other questions, please contact GlobalHealth Customer Care at 1-844-280-5555 or, for TTY users, 711 8:00 a.m. to 8:00 p.m., 7 days a week from October 1 – March 31 and 8:00 a.m. to 8:00 p.m., Monday – Friday, April 1 – September 30 or visit www.GlobalHealth.com/medicare.